

Case report

Nephrogenic Adenoma Of The Bladder.

Nunzio Di Naro¹, Marta Meloni¹, Tolu Giovanni Antonio^{1,2}.

1. Pathological Anatomy Operations Unit, "San Martino" Hospital, Oristano, Italy.

2. Director of Pathological Anatomy Operations Unit, "San Martino" Hospital, Oristano, Italy.

Abstract

We describe the clinical case of a older woman with a previous diagnosis of moderately and poorly differentiated and non-infiltrating bladder cancer (G2-G3) trated with Bacillus Calmette-Guerin (BCG) and with a subsequent histological diagnosis of nephrogenic adenoma. The material sent for histological examination, received in various fragments papillary-looking and fixed in 4% formalin, was sampled and included in paraffin. At the microscope in one of the fragments there was a cell proliferation with tubular architecture, consisting of monostratified cuboid cells with a large nucleus and scarce eosinophilic cytoplasm. These cells showed positivity for CK7 and PAX8, negativity for CK20 and focal positivity for GATA3, Ki67 showed a low proliferative index. The histopathological diagnosis was nephrogenic adenoma.

The subsequent follow-up of the patient, who is still alive and in good health, was negative. The differential diagnosis is with urothelial papilloma, low-grade urothelial neoplasms, prostatic adenocarcinoma, urothelial carcinoma of uncertain histology and clear cell bladder carcinoma, although the latter is rare.

Finally, since recurrence is frequent, constant follow-up is required.

Keywords : Urinary Bladder[1] , Nephrogenic adenoma[2], Nephrogenic Metaplasia[3] , Urinary Bladder neoplasms[4], Urologic Diseases[5], Urothelial Carcinoma[6].

CASE REPORT

A older woman, with a previous diagnosis of non-infiltrating, moderate-slightly differentiated urothelial carcinoma (G2-G3) treated with BCG, during subsequent ultrasound checks showed a neoformation of vegetative aspect of about 1x1 cm in the cavity bladder for which it was necessary to revise by means of cystoscopy and to remove it.

The material sent for histological examination, received in various fragments papillary-looking and fixed in 4% formalin, was sampled and included in paraffin.

At the microscope, in one of the fragments there was a cell proliferation with tubular architecture, consisting of monostratified cuboid cells with a large nucleus and scarce eosinophilic cytoplasm (fig 1,2,3).

On immunohistochemistry, these cells showed positivity for CK7 (fig 4) and PAX8 (fig 5,6), negativity for CK20 (fig 7) and focal positivity for GATA3 (fig 8).

Ki67 showed a low proliferative index (fig 9).

The histopathological diagnosis was nephrogenic adenoma.


The subsequent follow-up of the patient, who is still alive and in good health, was negative.

***Corresponding Author:** Marta Meloni, Pathological Anatomy Operations Unit, "San Martino" Hospital, Oristano, Italy. **Email:** martameloni1981@libero.it, **Orcid :** <https://orcid.org/0000-0003-4100-9848>

Received: 17-Feb-2025, Manuscript No. AOU-4548 ; **Editor Assigned:** 18-Feb-2025 ; **Reviewed:** 08-Mar-2025, QC No.AOU-4548 ; **Published:** 21-Mar-2025, **DOI:** 10.52338/aou.2025.4548

Citation: Marta Meloni. Nephrogenic adenoma of the bladder. Annals of Urology. 2025 March;10(1). doi: 10.52338/aou.2025.4548.

Copyright © 2025 Marta Meloni. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Figure 1. E.e. And Immunohistochemistry.

DISCUSSION

First described in 1949 (1), nephrogenic adenoma (or nephrogenic metaplasia) is considered a benign neoplasm or a metaplastic response of the bladder epithelium to chronic infection, stones or prolonged catheterisation (2); it is often diagnosed in patients who have undergone renal transplantation or who are undergoing BCG therapy for urothelial bladder cancer.(3)

On cystoscopy, it appears as low-grade urothelial carcinoma or as chronic cystitis, and at histological examination it is also necessary to evaluate the multiple aspects that the lesion can assume.

Macroscopically it is a sessile lesion, papillary or polypoid type, single, sometimes multiple.

Microscopically it is a heterogeneous lesion, showing a variety of aspects: tubular, papillary, fibromyxoid, flat, diffuse to large cells with clear cytoplasm (4,5,6).

Frequent occurrence of myxoid material surrounding the tubules, which may show bluish intraluminal myxoid inclusions mimicking signet ring cells.

Sometimes the neoplasm can invade the underlying muscle layer, so the possibility of malignant evolution cannot be excluded, especially in small biopsies (7).

The epithelium is cuboidal and may sometimes show nuclear atypia, with large nuclei and prominent nucleoli, with focal and superficial invasion of the muscle layer that may make difficult the differential diagnosis with a malignant carcinoma. Immunohistochemistry is positive for PAX-8, PAX-2, CK7, AMACR, EMA, usually negative for p63 (8,9,10).

The differential diagnosis is with urothelial papilloma, low-

grade urothelial neoplasms, prostatic adenocarcinoma, urothelial carcinoma of uncertain histology (11) and clear cell bladder carcinoma, although the latter is rare (12).

Finally, since recurrence is frequent, constant follow-up is required.

CONCLUSION

The aim of our case report is to broaden pathologists' knowledge of the histological variability of nephrogenic adenoma, which is a common diagnosis in urological pathology, but can be difficult to interpret under the microscope due to its extensive polymorphism.

Although most of the time nephrogenic adenoma presents with typical tubules and papillae, it may present with a varied spectrum of atypical histological features that may lead to a diagnostic error of malignancy.

Therefore, the pathologist must keep in mind the great variability of histological presentations of nephrogenic adenoma when investigating bladder or urinary tract lesions that are difficult to interpret (13,14,15) and keep it in the range of differential diagnostic possibilities, especially in patients treated with BCG or with a long history of bladder cancer recurrence, in order to make a correct diagnosis.

Availability Of Data

The data supporting the results of this study are available from the corresponding author upon request.

Conflicts Of Interest

The authors declare that there is no conflict of interest regarding the publication of this study.

REFERENCES

1. Davis TA: Hamartoma of the urinary bladder. *Northwest Med*, 1949;48:182-185
2. Ford TF, Watson GM, Cameron KM. Adenomatous Metaplasia (nephrogenic adenoma) of urothelium. An analysis of 70 cases. *Br J Urol* 1985;57(4):427-433
3. Stilment MM, Sivosky MB. Nephrogenic metaplasia associated with intravesical BCG treatment: report of 10 cases. *J Urol*. 1988;139 (1): 45-47
4. Oliva E, Young RH. Nephrogenic adenoma of the urinary tract: a review of the microscopic appearance of 80 cases with emphasis on unusual features. *Mod Pathol* 1995;8(7):722-730
5. Hansel DE, Nadasdy T, Epstein JI. Fibromyxoid nephrogenic adenoma: a newly recognized variant mimicking mucinous adenocarcinoma *Am J Surg Pathol* 2007;31(8):1231-1237
6. Pina-Oviedo S, Shen SS, Truong LD, et al. Flat pattern of nephrogenic adenoma: previously unrecognized pattern unveiled using PAX2 and PAX8 immunohistochemistry. *Mod Pathol* 2013;26:792-798
7. Alexiev BA, LeVeau CM: Nephrogenic adenoma of the urinary tract: a review *Int J Surg Pathol*, 2012;20(2):123-131
8. Xiao GQ, Burstein DE, Miller LK, et al. Nephrogenic adenoma Immunohistochemical evaluation for its etiology and differentiation from prostatic adenocarcinoma *Arch Pathol Lab Med*. 2006;130:805-810
9. Skinnider BF, Olivia E, Young RH, et al. Expression of P504S in nephrogenic adenoma: a significant immunohistochemical pitfall compounding differential diagnosis with prostatic adenocarcinoma *Am J Surg Pathol*
10. Isimbaldi G, Di Nuovo F, Sironi M, Catania G, Bonacina R, Privitera O, Spinelli M. Nephrogenic adenoma of the bladder. Morphological and immunophenotypic study with particular attention to differential diagnosis]. *Pathologica*. 1999 Jun;91(3):192-7. Italian. PMID: 10536465.
11. Lakshmi P. Kunju Nephrogenic Adenoma: report of a case and review of Morphologic Mimics *Arch Pathol Lab Med* 2010;134:1455-1459
12. Young RH, Scully RE: Nephrogenic adenoma. A report of 15 cases, review of the literature, and comparison with clear cell adenocarcinoma of the urinary tract. *Am J Surg Pathol*, 1986; 10:268-275
13. Amin W, Parwani AV. Nephrogenic adenoma. *Pathol Res Pract*. 2010 Oct 15;206(10):659-62. doi: 10.1016/j.prp.2010.06.001. Epub 2010 Jul 8. PMID: 20619548.
14. Santi R, Angulo JC, Nesi G, de Petris G, Kuroda N, Hes O, López JI. Common and uncommon features of nephrogenic adenoma revisited. *Pathol Res Pract*. 2019 Oct;215(10):152561. doi: 10.1016/j.prp.2019.152561. Epub 2019 Jul 25. PMID: 31358481.
15. Venyo AK. Nephrogenic Adenoma of the Urinary Bladder: A Review of the Literature. *Int Sch Res Notices*. 2015 Feb 2;2015:704982. doi: 10.1155/2015/704982. PMID: 27347540; PMCID: PMC4897129.