

Anxiety and depression when handicapped person stroke

Diham Zahi, Saila Mahir, Terman Azanmasso, Zatima Lmidmani

Correspondence to Author:

Diham Zahi, Department of Physical Medicine & Rehabilitation, Chu ibn rochd Casablanca, Morocco,

E-mail address: drzahidiham@gmail.com

Received Date: Dec 13, 2020

Accepted Date: Dec 16, 2020;

Published Date: January 18, 2021

Abstract

Objective: medical specialty sequelae of stroke square measure common, with depression and anxiety being the two most frequent manifestations.1–3 Our objective was to analyze the incidence of depression and anxiety in our patients once stroke.

Material/Patients and methods: A longitudinal study over twelve months from Gregorian calendar month 2014 to Gregorian calendar month 2015. It coated all cases of latest stroke supported the MPR service throughout this era. we tend to used HAD depression (Hospital Anxiety and Depression Scale Sigmond and Snaith).

Results: Of sixty four handicapped person patients followed in our department solely eighteen patients incontestible post-stroke depression (26.5%) and fourteen patients a state of tension (20.6%). The mean age was fifty seven.14 years with a variety from eighteen to eighty seven years. The sex- magnitude relation was one.1 for girls. the bulk of strokes were anaemia. medicine treatment was instituted for the eighteen patients. Conclusion: Depression ought to be sought-after consistently all told patients once stroke, particularly within the early part and at regular interviews with family. It compromises practical prognosis and it's a supply of demotivation particularly in rehabilitation.4–7

Keywords: depression, stroke, hemiplegia

Introduction

Neuropsychiatric manifestations of stroke (including affectional, cognitive, and behavioural symptoms) square measure even less seemingly to be recognized by patients or reportable within the literature.1 medical specialty sequelae of stroke square measure common, with depression and anxiety being the two most frequent manifestations.1–3 Recent proof estimates that one quarter of stroke patients develop anxiety,2 one third of stroke patients develop depression, and over 1/2 all stroke patients are laid low with depression at some purpose.3 Mood disorders considerably alter the lives of survivors and their families.4–7 It's best to spot them early and accurately so as to impose treatment.

Methods

In our study we tend to used Hospital Anxiety and Depression Scale (HADS)8–10 administered throughout a face-to-face interview a pair of, 4, six and twelve months once stroke, from Gregorian calendar month 2014 to Gregorian calendar month 2015 at the physical medication and rehabilitation department of IBN ROCHD University Hospitals.

The HADS consists of fourteen things and is split in 2 subscales directed at either depressive symptoms (HADS-D) or symptoms of tension (HADS-A) (7 things per subscale, vary 0-21 on every subscale). in step with the literature, each depression and anxiety is also outlined

by a HADS-subscale score >7.10 The HADS has incontestible smart psychological science properties, as well as smart internal consistency, in patients with stroke.11

All subjects had a clinical diagnosing of stroke that was created in step with the planet Health Organization observation. diagnosing confirmed with CT or resonance imaging proof of associate degree acute pathology, that was a part of the quality clinical stroke protocol. Patients were eligible if: 1) the diagnosing of cerebral stroke (ischemic or haemorrhagic) was clinically confirmed; 2) patients ought to be hemiplegic; 3) age was eighteen years or older; 4) comfortable information of the French language to complete the planned assessments. Patients with associate degree impaired level of consciousness, different acute medical or medical specialty malady, or a pre-morbid Axis I psychiatric diagnosing were excluded.

Clinical and medical histories assessed through meticulous medical specialty assessment and patient's interviews. Demographic data enclosed age, gender, legal status, living scenario, employment, and academic history. Concomitant medications and time since stroke were recorded.

Results

From associate degree initial cohort of one hundred thirty stroke patients, sixty four were enclosed. At one year post stroke, seventeen patients had died, fifteen patients had language difficulties, nine patients had a coffee mood, sixteen patients psychiatric issues apart from depression and nine patients were lost to follow- up. Mean age was fifty seven.14 years with a variety from eighteen to eighty seven years recent. Sex-ratio was one.1 for girls. the bulk of strokes were classified as anaemia (93%), with solely four haemorrhagic strokes (5%). mean solar time to the primary interview was twenty seven days with a variety from seven to fifty eight days. Of sixty four handicapped person patients followed in our department solely eighteen patients had post-stroke depression (26.5%) and fourteen patients a state of tension (20.6%) a pair of months once stroke. These results failed to extremely amendment four, six and twelve months later with eighteen cases of depression and sixteen cases of tension. medicine treatment (Fluoxetine twenty mg/ day) was instituted for the eighteen patients.

Discussion

The aim of our study was to explore the incidence for developing depressive symptoms and symptoms of tension once stroke. The known prevalence of twenty six.5% for depressive symptoms post stroke within the gift study was in line with the prevalence reportable in an exceedingly recent meta-analysis (33%, ninety fifth CI:23-43),3 whereas for symptoms of tension post stroke the determined prevalence during this study (20.6%) corresponded with the prevalence reportable in an exceedingly recent review (24%, ninety fifth CI:19-29).2

From a clinical perspective, our results indicate that the long-term risk of symptoms of depression and anxiety post stroke may be calculable supported a comparatively set of diagnostic data out there at 2 months post stroke. Psychological characteristics, early mood issues and neurosis, represent the core of those diagnostic sets. the first HADS-D makes an outsized contribution to work out chance of developing depressive symptoms and symptoms of tension post stroke.12 Previous studies13,14 found that stroke severity, in terms of practical dependency, makes a novel contribution to the chance of developing post stroke depressive symptoms.

Our study has limitations. It appears plausible that patients lost at follow up might have had a larger prevalence of mood disorders. Patients refusing assessments is also a lot of seemingly to possess mood disorders, then our results square measure so seemingly to underestimate actuality burden of symptoms of mood disorders. Moreover, the shortage of a full imaging assessment for different vital factors contributive to mood disorders (e.g., substantia alba changes and silent infarcts) might limit the generalization of our results. Another limit is that our assessment of symptoms was supported a metric “screening” scale, and not on a comprehensive clinical analysis. we tend to square measure aware that a brief scale isn’t a substitute for professional clinical assessment; but, the dimensions we tend to used is one in every of the foremost wide used tools to find mood disorders once stroke.¹⁵

Conclusion

The results of this study counsel that depression and anxiety were usually practised post stroke. Post-stroke mood disorders ought to be sought-after consistently all told handicapped person patients, particularly within the early part and at regular interviews with family. They compromise practical prognosis and that they square measure a supply of demotivation particularly in rehabilitation.

Conflict of interest

The author declares no conflict of interest.

References

1. Hackett ML, Köhler S, O'Brien JT, Mead GE. Neuropsychiatric outcomes of stroke. *Lancet Neurol*. 2014;13(5):525–534.
2. Campbell Burton CA, Murray J, Holmes J, et al. Frequency of anxiety after stroke: a systematic review and meta-analysis of observational studies. *Int J Stroke*. 2013;8(7):545–559.
3. Ayerbe L, Ayis S, Crichton S. The natural history of depression up to 15 years after stroke: the South London Stroke Register. *Stroke*. 2013;44(4):1105–1110.
4. Nelson LD, Cicchetti D, Satz P. Emotional sequelae of stroke—a longitudinal perspective. *J Clin Exp Neuropsychol*. 1994;16(5):796–806.
5. Parikh RM, Robinson RG, Lipsey JR, et al. The impact of post stroke depression on recovery in activities of daily living over a 2-year follow-up. *Arch Neurol*. 1990;47(7):785–789.
6. Morrison V, Pollard B, Johnston M. Anxiety and depression 3 years following stroke: demographic, clinical, and psychological predictors. *J Psychosom Res*. 2005;59(4):209–213.
7. Flick CL. Stroke rehabilitation. 4. Stroke outcome and psychosocial consequences. *Arch Phys Med Rehabil*. 1999;80(5 Suppl 1):S21–S26.
8. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand*. 1983;67(6):361–370.
9. Brennan C, Worrall-Davies A, McMillan D. The hospital anxiety and depression scale: A diagnostic meta-analysis of case-finding ability. *J Psychosom Res*. 2010;69(4):371–378.
10. Aben I, Verhey F, Lousberg R, et al. Honig Validity of the beck depression inventory, hospital anxiety and depression scale, SCL-90, and hamilton depression rating scale as screening instruments for depression in stroke patients. *Psychosomatics*. 2002;43(5):386–393.
11. Spinhoven P, Ormel J, Sloekers PP, et al. A validation study of the hospital anxiety and depression scale (hads) in different groups of dutch subjects. *Psychol Med*. 1997;27(2):363–370.
12. Kootker JA, van Mierlo ML, Hendriks JC, et al. Risk factors for symptoms of depression and anxiety one year post stroke: a longitudinal study. *Archives of physical medicine and rehabilitation*. 2016;97(6):919–928.
13. De Ryck A, Brouns R, Geurden M, et al. (2014) Risk factors for poststroke depression: Identification of inconsistencies based on a systematic review. *J Geriatr Psychiatry Neurol*. 2014;27(3):147–158.
14. Kutlubaev MA, Hackett ML. Predictors of depression after stroke and impact of depression on stroke outcome: An updated systematic review of observational studies. *Int J Stroke*. 2014;9(8):1026–1036.
15. Lees R, Fearon P, Harrison JK, Quinn TJ. Cognitive and mood assessment in stroke research: focused review of contemporary studies. *Stroke*. 2012;43(6):1678–1680.