Review Article

Annals Of Agricultural Science And Technology

Soil Application Of Plant Growth-Promoting Fungi: A Sustainable Strategy For Agriculture.

Muskan Bhullar[1]

[1]*PG Department of Agriculture, Division of Plant pathology Khalsa College Amritsar 143001*

Corresponding author
Dr. Muskan Bhullar ,
PG Department of Agriculture, Division of Plant pathology Khalsa College Amritsar 143001.
Email : msknbhullar@gmail.com

Received Date : November 15, 2024
Accepted Date : November 16, 2024
Published Date : December 18, 2024

Copyright *© 2024 Dr. Muskan Bhullar. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.*

ABSTRACT

PGPF (Plant Growth-Promoting Fungi) application is a revolutionary sustainable approach that enhances crop yields, improves soil health, and mitigates environmental degradation. PGPF interacts with plants, stimulating root growth, and boosting nutrient uptake, leading to improved crop productivity and quality. The fungal community also induces systemic resistance in plants through the expression of defence-related enzymes (such as peroxidase, polyphenol oxidase, and chalcone synthase) and defence chemicals (such as phytoalexin and anti-microbial phenolic compounds) for conferring structural and chemical barrier against pathogens and pests. Additionally, PGPF promotes soil biodiversity, structure, and fertility, increasing water retention and aeration. This approach reduces the reliance on chemical fertilizers and pesticides, minimizing soil pollution and environmental harm. By adopting PGPF application, farmers can achieve sustainable agriculture, ensuring food security and environmental conservation.

Keywords
Mycoparasitism, Phytohormones, Plant Growth-Promoting Fungi, Rhizosphere Sustainable Agriculture, Systemic Resistance.

INTRODUCTION

PGPF are a heterogeneous group of soil-dwelling, non-pathogenic saprophytic fungi that establish a close association with plants and promote plant growth and health through several activities (Naziya *et al.*, 2020). Further, fungi under PGPF may differ distinctly from each other with respect to their taxonomy, habitats, physiology, and even to their interactions with plants. The term PGPF is not absolute, rather it is an operational term (Bent, 2006), as all fungi that promote plant growth are not PGPF, for example, mycorrhizal fungi, which are known to boost the growth of the plants, are not considered as PGPF. An important feature that gives PGPF a different identity is that the they are non-symbiotic saprotrophic fungi that live freely on a zone of soil at the vicinity of the root or the interior of the root itself, whereas mycorrhizal fungi behave as obligate biotrophs and develop an intimate association with the roots of most host plants (Hossain *et al*., 2017a, b). Root colonization ability is considered as one of the most important characteristics of PGPF which helps to promote plant growth (Islam *et al*., 2014). Fungi of the genera such as *Aspergillus, Fusarium, Penicillium, Piriformospora, Phytophthora, Rhizoctonia, Phoma*, and *Trichoderma* are the strains mostly used in research as PGPF (Hossain *et al*., 2017a, b; Javaid et al., 2020; Murali et al., 2021). The non-pathogenic fungi such as *Pythium oligandrum* and *Phytophthora cryptogea* colonizing the root ecosystems are also considered as PGPF (Benhamou *et al*., 2012; Bent 2006). PGP fungi have reported from different genera of phyla Chytridiomycota, Zygomycota, Glomeromycota, Ascomycota, and Basidiomycota. The beneficial effects of Plant Growth-Promoting Fungi (PGPF) are linked to their ability to colonize roots, produce growth hormones, facilitate mineralization, enhance nutrient uptake, control diseases through antagonistic mechanisms, and trigger defense strategies against pathogens. These defense strategies include inducing systemic resistance (ISR) and systemic acquired resistance (SAR) in plants, which involves the production of defense enzymes, chemicals, and pathogenesis-related proteins (PR-proteins) (Islam *et al*., 2014; Nogueira-Lopez *et al.*, 2020). The beneficial fungi promote the plant via directed multifarious plant growth-promoting (PGP) attributes like micronutrients solubilization (phosphorus, potassium and zinc) and production of plant growth regulators like auxin, gibberellins, cytokinin and ethylene or indirectly via the antagonistic substances, production of siderophores, antibiotic and synthesis of cell wall lysing enzymes like

cellulases, glycosidase and gluconase (Abo Nouh, 2019; Urja and Meenu, 2010). By harnessing the benefits of beneficial soil microorganisms, including fungi, soil health can be substantially enhanced, diseases can be controlled through antagonistic interactions and induced systemic resistance, plant growth can be promoted, and a more sustainable alternative to synthetic chemical fertilizers can be provided (Bhardwaj *et al*., 2014). These prospective PGPF would play an important role in agriculture for long-term productivity, soil health management, and environmental restoration as a cost-effective input in the next decades, perhaps providing substantial relief for food security.

Rhizosphere

Rhizosphere is the soil zone, which is influenced by the roots. Plants share a micro-ecosystem at the vicinity of the plant root system comprising hot spot zone of the microbial community (like bacteria, fungi, nematodes, viruses, arthropods, oomycetes, protozoa, algae, and archaea), of which bacteria and fungi are most common and extensively studied (Akinola and Babalola 2021). The plant rhizosphere harbours both beneficial and pathogenic microorganisms comprising up to 1011 microbes per gram of soil and above 3000 prokaryotic species in general; and therefore, it represents a composite ecosystem on earth (Hossain et al., 2017a, b; Mendes et al., 2013). This rhizospheric zone offers great opportunities for plant–microbial interactions, and therefore significantly affects plant growth, disease resistance, and nutrient recycling (Akinola and Babalola, 2021). Understanding the complex microbial interactions in the rhizosphere is essential for developing organic farming practices that reduce reliance on synthetic chemical fertilizers, which can harm the environment and surrounding ecosystems (Rascovan et al., 2016).

1.1 Plant Growth Promotional Activities of PGPF

The plant growth promotional activities of PGPF are attributed to the production of plant growth hormone and mineralization and as such many more. Research has consistently shown that PGPF play a significant role in enhancing various aspects of plant growth and development, including seed germination, seedling vigor, shoot and root growth, photosynthetic efficiency, flowering, and ultimately, crop yield (Hossain and Sultana, 2020)

Production of Plant Growth Hormone

Plant growth hormones also called phytohormones help to regulate the growth of the plants through various developmental processes. Phytohormones, specifically auxins (IAA), gibberellins, and cytokinin can exogenously produce by Plant Growth-Promoting Fungi (PGPF) and play a pivotal role in regulating plant growth and development. IAA and gibberellins are particularly significant, as they induce crucial physiological responses during various stages of plant growth (Islam et al., 2014). The production of IAA, a highly important and widely distributed phytohormone, has been reported in several fungi, including Trichoderma, Penicillium, Aspergillus, Fusarium, Talaromyces, and Mortierella, in host plants such as chickpea, rice, and wheat, resulting in enhanced growth and yield (Abri et al., 2015; Kumar et al., 2017; Murali et al., 2021). Gibberellic acid (GA) is a phytohormone of significance, produced by various fungi, including Fusarium, Aspergillus, and Penicillium, which contributes substantially to plant growth and developmental processes, in addition to conferring tolerance to abiotic stress (Syamsia et al., 2021). Notably, the GA produced by Cladosporium species in wheat and cucumber plants has been demonstrated to enhance plant growth (Hamayun et al., 2010). Moreover, endophytic fungi, such as Penicillium citrinum and Aspergillus fumigatus, have been reported to promote plant growth by secreting GAs in the rhizosphere, thereby stimulating plant development and growth (Ahmad et al., 2010). Another important plant growth regulator, cytokinin (predominantly zeatin), elicited by Piriformospora spp., Phoma spp., and Trichoderma spp. caused growth promotion in melon and Arabidopsis (Martínez-Medina et al., 2014; (Hossain and Sultana 2020; Speakman and Kruger 1984).

Mineralization

Mineral availability in the rhizosphere is controlled by combined effects of soil properties, plant characteristics, and root-microorganism interactions (Jones et al., 2004; Rengel and Marschner, 2005). Rhizospheric fungi can enhance mineral uptake and availability, compensating for deficiencies through their symbiotic relationship with plant roots.

Phosphorous Solubilization

Plant Growth-Promoting Fungi (PGPF) in the rhizosphere significantly increase phosphorus (P) availability for plants by solubilizing phosphate compounds. Research has shown that Phosphate-Solubilizing Fungi (PSF) can convert insoluble phosphate into soluble forms, providing a promising alternative to phosphorus fertilizers (Alam et al., 2002; Chabot et al., 1996; Pal 1998). The fungi achieve this by producing various organic acids like tartaric acid, succinic acid, oxalic acid, malic acid, 2-ketogluconic acid, glyoxylic acid, gluconic acid, fumaric acid, citric acid and alpha-ketobutyric acid and enzyme phosphatase (Devi et al., 2020). All the release compounds solubilize the phosphorus and avail the soluble inorganic form that can be assimilated by the plants. A number of fungal

species including P. bilaji, Penicillium spp. (Patil et al. 2012), P. oxalicum (Li et al. 2016a) Aspergillus niger, Penicillium notatum (Din et al. 2019; Malviya et al. 2011), Aspergillus awamori (Jain et al. 2012a), Penicillium bilaii (Ram et al. 2015), Trichosporon beigelii, Rhodotrula aurantiaca, Cryptococcus luteolus, Zygoascus hellenicus, P. purpurogenum var. rubrisclerotium, Neosartorya fisheri, and Candida montana (Gizaw et al. 2017), Talaromyces aurantiacus, Aspergillus neoniger (Zhang et al. 2018a), and Trichoderma spp. (Bononi et al. 2020) have been reported for the solubilization of phosphorus. Rhizospheric fungi Penicillium, Aspergillus, Trichoderma, Phoma, Rhizoctonia, Rhizopus, and Alternaria have been documented for their efficiency in solubilizing the insoluble phosphate (Alori et al., 2017; Dotaniya and Meena 2015). For the potassium solubilizination we can use large amount of microorganism like as Aspergillus spp., Agrobacterium tumefaciens, B. pumilus, B. subtilis, B. circulans, B. edaphicus, B. mucilaginosus, Flavobacterium spp. and Rhizobium spp. (Gundala et al. 2013; Keshavarz Zarjani et al. 2013; Maurya et al. 2015; Meena et al. 2014a; Meena et al. 2014b). This potash- solubilizing biofertilizer can be applied in crop production and yield combination with potassium solubilizing microbiome are Azospririllium, Azotobacter, Azospirillum, Acetobacter and Rhizobium (Bahadur et al. 2016). Among the PGPF, different Trichoderma strains have been exploited to enhance mineralization and mineral absorption of Fe, N, P, and K, and increase the accessibility of ammonium, nitrogen, zinc, copper, iron, and manganese (Molla et al., 2012).

1.2 Management of Disease

The use of Plant Growth-Promoting Fungi (PGPF) in plants is environmentally friendly approach to manage diseases, which triggers a long-lasting activation of innate immunity of plants. Apart from promoting plant growth, PGPF employs various strategies to protect plants from pathogens by inducing defence resistance. The primary mechanism of PGPF in sustainable disease management involves colonizing plant roots, facilitating nutrient uptake, and stimulating plant growth (Hossain et al., 2017a, b; Murali et al., 2013). The key disease management strategies employed by PGPF can be summarized as follows:

Antagonism

The bio-control mechanism against disease-causing pathogens can be achieved through the antagonistic efficacy of microorganisms. PGPF exerts its bio-control mechanism through multiple strategies, including antibiosis, competition, and parasitism. Antibiosis occurs through the production of antibiotics and biosurfactants, which inhibit pathogen growth. Competition ensues as PGPF competes with pathogens for colonization sites, nutrients, and minerals. Additionally, PGPF produces extracellular cell wall-degrading enzymes, such as chitinase and β-1,3-glucanase, which break down pathogen cell walls, ultimately reducing damage to plants (Berg et al., 2005). Different PGPF, namely, Trichoderma, Gliocladium virens, Phoma sp., Fusarium equiseti, and Penicillium simplicissimum have been reported to be antagonistic against Rhizoctonia solani, Pythium aphanidermatum, Pythium irregulare, Sclerotium rolfsii, Fusarium oxysporum, Pseudomonas syringae, and Colletotrichum orbiculare (Lewis et al., 1998; Murali et al., 2021). Patale and Mukadam (2011) have successfully tested the antagonistic activity of Trichoderma viride and Trichoderma harzianum against seven pathogenic fungi, namely, Aspergillus niger, A. flavus, Phytophthora spp., Fusarium oxysporum, Rhizoctonia solani, Penicillium notatum, and Alternaria solani. Gliovirin, an antibiotic produced by Gliocladium virens, was shown to inhibit the growth of Pythium ultimum (Howell and Stipanovic 1983).


PGPF and induced resistance

Induced resistance is triggered by pathogens or insects, activating defensive compounds. It has two categories: Induced Systemic Resistance (ISR) and Systemic Acquired Resistance (SAR). ISR is activated through the jasmonate pathway (Cong and Lingyun, 2019), triggered by pathogens, drought, herbivores, and mechanical injuries. SAR is regulated by the salicylic acid-dependent signaling pathway, involving local and systemic increases in salicylic acid, leading to the promotion of pathogenesis-related proteins (Backer et al., 2019). PGPF can elevate plant's defence against insects and diseases by triggering resistance or activating natural defence responses (Adesemoye and Kloepper, 2009). The level of induced resistance in plants can vary depending on factors such as the source, type, and intensity of stimuli (Aranega-Bou et al., 2014). Trichoderma strains inoculated to the rhizosphere protect host plants against numerous pathogens including bacteria, and fungi, due to the induction of resistance responses similar to the hypersensitive mechanism, SAR and ISR in plants (Singh et al. 2016; Jyoti et al., 2014), as peroxidase reactive oxygen species (ROS) activity was triggered by T. virens in cotton plants reported by Singh et al. (2016).The use of non-pathogenic strains of F. oxysporum to control wilt disease has been detected for many crops (Ahmad et al., 2018). Pythium oligandrum has shown the ability to control soil-borne phytopathogens either in the laboratory or in the field. P. oligandrum oospores have been used as seed treatments that reduce damping-off disease in sugarbeet caused by P. ultimum (Rocha et al., 2019). Also, P. nunn is an antagonistic fungus or mycoparasite of pathogens such as Pythium ultimum, P. aphanidermatum P. vexans, Rhizoctonia

solani, Phytophthora parasitica, and P. cinnamomi. Moreover, Aspergillus and Penicillium species were effective against the white-rot disease caused by basidiomycetes (Kowalczyk et al., 2019).

Penicillium oxalicum, a plant growth promoting fungus (PGPF) isolated from pearl millet rhizosphere soil, exhibited a considerable increase in chitinase activity (Murali and Amruthesh 2015). Trichoderma atroviride TRS25 increased PPO and PAL enzyme activity when the cucumber plant was inoculated with Rhizoctonia solani (Nawrocka et al., 2018). Trichoderma spp. is widely studied for their role in controlling the phytopathogen by the production of cell wall degrading enzymes such as cellulases, chitinases, and glucanases (Nogueira-Lopez et al., 2020).

Figure 1: PGPF in contributing overall growth of the plant through direct and indirect mechanisms (Mandal P and Tiru Z, 2024).



Bioformulations

Biocontrol agents (bio-agent) comprising fungi have become attractive in terms of sustainable management of diseases and improved quality of crop productivity (Hussain et al., 2020). The antagonistic property of PGPF can be successfully exploited through proper identification of efficient bio-control agents, their multiplication, and formulation for delivery. A large number of bio-based products are being produced and sold worldwide in the form of granules, wettable powders, dusts, and aqueous or oil-based liquid products using different carriers to control fungal pathogens (Ardakani et al., 2009; Nega 2014). The application of Trichoderma-based bio-fertilizer (composted of cattle manure + inoculum) not only produced the antifungal compound which may suppress the pathogen but potentially improved grassland biomass (Zhang et al., 2018). Bio-organic fertilizers (BOFs) enriched with Trichoderma and animal manure have been found not only to cause plant growth promotion but also found to have the controlling effect against Fusarium wilt in cucumber plants (Chen et al., 2011; Zhang et al., 2013, 2016). The foliar sprays of the liquid formulation of Penicillium oxalicum (6 X 106 CFU ml−1) with sodium alginate (0.5%) and Tween 80 (0.01%) substantially improved the yield and acted as biofungicide for controlling mango malformation (Haggag and El Soud, 2013).

1.3 Abiotic Stress Management

Plant growth promoting fungi (PGPF) are recognised for alleviating a variety of abiotic challenges (heavy metal stress, water stress, temperature stress, and salt stress). Penicillium species isolated from groudnut rhizosphere soil were reported to improve salinity tolerance in sesame plants, as well as disease resistance and plant growth promotion (Radhakrishnan et al., 2014). Trichoderma harzianum promotes root growth and aids in water absorption and nutrient intake during osmotic stress. Distinct strain of Trichoderma has been carefully studied for its ability to mitigate oxidative, salinity, drought, and osmotic stress in plants (Zaidi et al., 2014). Microsphaeropsis, Mucor, Steganosporium, Phoma, Aspergillus, Alternaria, and Peyronellaea have been shown to protect Arabidopsis plants from heavy metal buildup (Murali et al., 2021). Trichoderma has been reported to assist plants increase the activity of antioxidant enzymes, hence protecting plants from ROS production and membrane damage in stressful conditions (Guler et al., 2016).

Conclusion and Future Prospects

Unlocking the benefits of Plant Growth-Promoting Fungi (PGPF) is crucial for developing innovative strategies to enhance crop yields and effectively manage crop diseases, leading to improved agricultural productivity and sustainability. Despite their potential, Plant Growth-Promoting Fungi (PGPF) face challenges in practical applications due to inconsistent performance, likely caused by genetic, environmental, and other factors. In order to popularize the widespread use of PGPF, the development of some innovative and effective techniques for their mass culture, formulation, and application of these fungi are urgently needed to be addressed. Advanced molecular tools and techniques can give more insight into mechanisms and outcomes of plant-microbial interactions. Additionally, biotechnological innovations, such as genetic modification and gene transfer, can enhance PGPF's benefits, leading to more promising and effective solutions for widespread adoption. PGPFs are successfully used in many countries, but some still rely on chemical fungicides. Addressing challenges and limitations is crucial to promote wider adoption and sustainable agriculture.

LITERATURE CITED

1. Abo Nouh, I (2019) Role of plant growth-promoting fungi in sustainable agriculture. Journal of Agricultural Science and Technology 19(2) :257-276.

2. Abri T K, Sengin E, Sjahrir R (2015) Production of indole acetic acid (IAA) hormone from fungal isolates collected from rhizosphere of aromatic rice in Tana Toraja. International Journal of Current Research in Biosciences and Plant Biology 2: 198–201

3. Adesemoye A O, Kloepper J W (2009) Plant-microbe interactions in enhanced fertilizer use efficiency. Applied Microbiology Biotechnology 85: 1–12.

4. Ahmad M, Pataczek L, Hilger T H, Zahir Z A (2018) Perspectives of Microbial Inoculation for Sustainable Development and Environmental Management. Frontiers in Microbiology 9: 2992.

5. Akinola S A and Babalola O O (2021) The fungal and archaeal community within plant rhizosphere: a review on their contribution to crop safety. Journal of Plant Nutrition 44: 600–618

6. Alam S, Khalil S, Ayub N, et al. (2002) In vitro solubilization of inorganic phosphate by phosphate solubilizing microorganisms (PSM) from maize rhizosphere. International Journal of Agricultural Biology 4(3): 454-458.

7. Alori E T, Glick B R, Babalola O O (2017) Microbial phosphorus solubilization and its potential for use in sustainable agriculture. Frontiers in Microbiology 8: 971.

8. Aranega-Bou P, de la O Leyva M, Finiti I, García-Agustín P, González-Bosch C (2014) Priming of plant resistance by natural compounds. Hexanoic acid as a model. Frontiers in Plant Science 5: 488.

9. Ardakani S S, Heydari A, Khorasani N A (2009) Preparation of new biofungicides using antagonistic bacteria and mineral compounds for controlling cotton seedling damping-off disease. Journal of Plant Protection Research 49: 1

10. Backer R, Naidoo S, den Berg N. 2019 – The Non Expressor of Pathogenesis-Related Genes 1) NPR1(and Related Family: Mechanistic Insights in Plant Disease Resistance. Frontiers in Plant Science 10: 102

11. Bahadur I, Maurya B R, Kumar A, Meena V S, Raghuwanshi R (2016) Towards the soil sustainability and potassium-solubilizing microorganisms. In: Meena VS, Maurya BR, Verma JP, Meena RS (eds) Potassium solubilizing microorganisms for sustainable agriculture. Springer India, New Delhi, pp 255-266

12. Benhamou N, Le Floch G, Vallance J (2012) Pythium oligandrum: an example of opportunistic success. Microbiology 158:2679–2694

13. Bent E (2006) Induced systemic resistance mediated by plant growth-promoting rhizobacteria (PGPR) and fungi (PGPF). In: Tuzun S, Bent E (eds) Multigenic and induced systemic resistance in plants. Springer, Boston, pp 225-258.

14. Berg G, Zachow C, Lottmann J et al (2005) Impact of plant species and site on rhizosphere-associated fungi antagonistic to Verticillium dahlia. Applied and Environmental Microbiology 71: 4203–4213

15. Bhardwaj D, Ansari M W, Sahoo R K and Tuteja N (2014) Biofertilizers: A sustainable alternative to chemical fertilizers for plant growth and soil health. Journal of Environmental Science and Health 49: 539-547

16. Bononi L, Chiaramonte J B, Pansa C C, Moitinho M A, Melo I S (2020) Phosphorus-solubilizing Trichoderma spp. from Amazon soils improve soybean plant growth. Scientific Report 10:1-13

17. Chabot R, Antoun H, Cescas MP (1996) Growth promotion of maize and lettuce by phosphate-solubilizing Rhizobium leguminosarum biovar. Phaseoli Plant and Soil 184(2):311-321.

18. Chen L, Yang X, Raza W (2011) Solid-state fermentation of agro-industrial wastes to produce bioorganic fertilizer for the biocontrol of Fusarium wilt of cucumber in continuously cropped soil. Bioresource Technology 102: 3900–3910

19. Cong Z, Lingyun Z (2019) An account of in silico identification tools of secreted effector proteins in bacteria and future challenges. Briefings in Bioinformatics 20: 110–129

20. Devi R, Kaur T, Kour D (2020) Beneficial fungal

communities from different habitats and their roles in plant growth promotion and soil health. Microbial Biosystems 5(1): 21-47.

21. Din M, Nelofer R, Salman M, Khan F H, Khan A, Ahmad M (2019) Production of nitrogen fixing Azotobacter (SR-4) and phosphorus solubilizing Aspergillus niger and their evaluation on Lagenaria siceraria and Abelmoschus esculentus. Biotechnol Rep 22: 00323

22. Dotaniya M L, Meena V (2015) Rhizosphere effect on nutrient availability in soil and its uptake by plants: a review. Proceedings of the National Academy of Sciences, India Section B: Biological Sciences 85(1):1-12.

23. Gizaw B, Tsegay Z, Tefera G, Aynalem E, Wassie M, Abatneh E (2017) Phosphate solubilizing fungi isolated and characterized from Teff rhizosphere soil collected from North Showa zone, Ethiopia. African Journal of Microbiological Research 11: 687-696

24. Guler N S, Pehlivan N, Karaoglu S A (2016) Trichoderma atroviride ID20G inoculation ameliorates drought stress-induced damages by improving antioxidant defence in maize seedlings. Acta Physiologiae Plantarum 38(10): 1-12.

25. Gundala P, Chinthala P, Sreenivasulu B (2013) A new facultative alkaliphilic, potassium solubilizing, Bacillus Sp. SVUNM9 isolated from mica cores of Nellore District, Andhra Pradesh, India. Research and Reviews. Journal of Microbiology and Biotechnology 2:1-7

26. Haggag W M and El Soud M A (2013) Pilot-scale production and optimizing of cellulolytic Penicillium oxalicum for controlling of mango malformation. Agricultural Science 4:165–17

27. Hamayun M, Khan S A, Khan A L (2010) Gibberellin production and plant growth promotion from pure cultures of Cladosporium sp. MH-6 isolated from cucumber (Cucumis sativus L.). Mycologia 102(5): 989-995

28. Hossain M M, Sultana F (2020) Application and mechanisms of plant growth promoting fungi (PGPF) for phytostimulation. In Das SK (ed) Organic Agriculture, IntechOpen, London/UK, pp 65-94.

29. Hossain MM, Sultana F, Hyakumachi M (2017a) Role of ethylene signalling in growth and systemic resistance induction by the plant growth-promoting fungus Penicillium viridicatum in Arabidopsis. Journal of Phytopathology 165: 432–441

30. Hossain M M, Sultana F, Islam S (2017b) Plant growth-promoting fungi (PGPF): phytostimulation and induced systemic resistance. In Singh D, Singh H, Prabha R (eds) Plant-microbe interactions in agro-ecological perspectives, Springer, Singapore, pp 135-191.

31. Howell C R and Stipanovic R D (1983) Gliovirin, a new antibiotic from Gliocladium virens, and its role in the biological control of Pythium ultimum. Canadian Journal of Microbiology 29: 321–324

32. Hussain T, Akthar N, Aminedi R (2020) Role of the potent microbial based bioagents and their emerging strategies for the eco-friendly management of agricultural phytopathogens. In Singh J, Yadav AN (eds) Natural bioactive products in sustainable agriculture. Springer, Singapore, pp 45–66

33. Islam S, Akanda A M, Sultana F (2014) Chilli rhizosphere fungus Aspergillus spp. PPA1 promotes vegetative growth of cucumber (Cucumis sativus) plants upon root colonisation. Archives of Phytopathology and Plant Protection 47(11): 1231-1238.

34. Javaid A, Afzal R, Shoaib A (2020) Biological management of southern blight of chili by Penicillium oxalicum and leaves of Eucalyptus citriodora International Journal of Agricultural Biology 23: 93-102

35. Jones D L, Hodge A, Kuzyakov Y (2004) Plant and mycorrhizal regulation of rhizodeposition. New Phytologist 163(3):459-480

36. Jyoti A, Singh AK, Dubey M, Kumar S (2014) Interaction of inducible nitric oxide synthase with rac2 regulates reactive oxygen and nitrogen species generation in the human neutrophil phagosomes: implication in microbial killing. Antioxidants and Redox Signaling 20: 417–431.

37. Keshavarz Zarjani J, Aliasgharzad N, Oustan S, Emadi M, Ahmadi A (2013) Isolation and characterization of potassium solubilizing bacteria in some Iranian soils. Archives of Agronomy and Soil Science 59:1713-1723

38. Kowalczyk J E, Peng M, Pawlowski M, Lipzen A (2019)

The White-Rot Basidiomycete Dichomitus squalens Shows Highly Specific Transcriptional Response to Lignocellulose Related Aromatic Compounds. Frontiers in Bioengineering and Biotechnology 7: 229.

39. Kumar NV, Rajam KS, Rani ME (2017) Plant growth promotion efficacy of indole acetic acid (IAA) produced by a mangrove associated fungi-Trichoderma viride VKF3. International Journal of Current Microbiology and Applied Sciences 6(10): 2692-2701.

40. Lewis J, Larkin R, Rogers D (1998) A formulation of Trichoderma and Gliocladium to reduce damping-off caused by Rhizoctonia solani and saprophytic growth of the pathogen in soilless mix. Plant Disease 82: 501–506

41. Li Y, Wang Z, Liu X, Song Z (2016)– Siderophore Biosynthesis but Not Reductive Iron Assimilation Is Essential for the Dimorphic Fungus Nomuraea rileyi Conidiation, Dimorphism Transition, Resistance to Oxidative Stress, Pigmented Microsclerotium Formation, and Virulence. Frontiers in Microbiology 7: 931.

42. Malviya J, Singh K, Joshi V (2011) Effect of phosphate solubilizing fungi on growth and nutrient uptake of ground nut) Arachis hypogaea (plants. Advanced Biomedical Research 2: 110– 113.

43. Mandal P and Tiru Z (2022) Soil application of plant growth promoting fungi for sustainable agriculture in the new decade. In Plant Stress: Challenges and Management in the New Decade (pp. 321-330). Cham: Springer International Publishing.

44. Martínez-Medina A, Alguacil M D M, Pascual J Al (2014) Phytohormone profiles induced by Trichoderma isolates correspond with their biocontrol and plant growth-promoting activity on melon plants. Journal of Chemical Ecology 40:804–815

45. Masunaka A, Hyakumachi M, Takenaka S (2009) Plant growth-promoting fungus, Trichoderma koningi suppresses isoflavonoid phytoalexin vestitol production for colonization on/in the roots of Lotus japonicus. Microbes and Environments 26(2):128-134.

46. Maurya B R, Kumar A, Raghuwanshi R, Bahadur I, Meena V (2015) Effect of phosphate solubilizing isolates on growth, yield and phosphate acquisition by rice and wheat crops. African Journal of Microbiological Research 9:1367- 1375

47. Meena V, Maurya B, Bahadur I (2014a) Potassium solubilization by bacterial strain in waste mica. Bangladesh Journal of Botany 43: 235-237

48. Meena VS, Maurya B, Verma JP (2014b) Does a rhizospheric microorganism enhance K+ availability in agricultural soils? Microbiological Research 169: 337-347

49. Mendes R, Garbeva P, Raaijmakers J M (2013) The rhizosphere microbiome: significance of plant beneficial, plant pathogenic, and human pathogenic microorganisms. Federation of European Microbiological Societies 37: 634–663

50. Molla A H, Manjurul Haque M, Amdadul Haque M (2012) Trichoderma-enriched biofertilizer enhances production and nutritional quality of tomato (Lycopersicon esculentum Mill.) and minimizes NPK fertilizer use. Agricultural Research 1(3): 265-272.

51. Murali M and Amruthesh KN (2015) Plant growth-promoting fungus Penicillium oxalicum enhances plant growth and induces resistance in pearl millet against downy mildew disease. Journal of Phytopathology 163:743–754

52. Murali M, Naziya B, Ansari M A (2021) Bioprospecting of rhizosphere-resident fungi: their role and importance in sustainable agriculture. Journal of Fungi 7(3): 314.

53. Murali M, Sudisha J, Amruthesh K (2013) Rhizosphere fungus Penicillium chrysogenum promotes growth and induces defence-related genes and downy mildew disease resistance in pearl millet. Plant Biology 15: 111–118

54. Naziya B, Murali M, Amruthesh K N (2020) Plant growth-promoting fungi (PGPF) instigate plant growth and induce disease resistance in Capsicum annuum L. upon infection with Colletotrichum capsici (Syd.) Butler & Bisby. Biomolecules 10(1): 41

55. Nega A (2014) Review on concepts in biological control of plant pathogens. Journal of Biological Agriculture and Healthcare 4: 33–54

56. Nogueira-Lopez G, Lawry R, Echaide-Aquino JF (2020)

Chemical communication between Trichoderma and plants. In Gupta VK, Zeilinger S, Singh HB, Druzhinina Ib (eds) New and future developments in microbial biotechnology and bioengineering. Elsevier, pp 109–139

57. Pal SS (1998) Interactions of an acid-tolerant strain of phosphate-solubilizing bacteria with a few acid-tolerant crops. Plant and Soil 198(1):169-177.

58. Patale S and Mukadam D (2011) Management of plant pathogenic fungi by using Trichoderma species. Bioscience and Disease Journal 2:36–37

59. Patil PM, Kuligod V, Hebsur N, Patil C, Kulkarni G (2012) Effect of phosphate solubilizing fungi and phosphorus levels on growth, yield and nutrient content in maize (Zea mays). Karnataka Journal of Agricultural Science 25

60. Radhakrishnan R, Kang S M, Baek I Y (2014) Characterization of plant growth-promoting traits of Penicillium species against the effects of high soil salinity and root disease. Journal of Plant Interactions 9(1):754-762.

61. Ram H, Malik S, Dhaliwal S, Kumar B, Singh Y (2015) Growth and productivity of wheat affected by phosphorus-solubilizing fungi and phosphorus levels. Plant Soil Environment 61:122-126

62. Rascovan N, Carbonetto B, Perrig D et al (2016) Integrated analysis of root microbiomes of soybean and wheat from agricultural fields. Scientific Reports 6:1–12

63. Rengel Z and Marschner P (2005) Nutrient availability and management in the rhizosphere: exploiting genotypic differences. New Phytologist 168(2): 305-312

64. Rocha I, Ma Y, Souza-Alonso P, Vosátka M (2019) Seed Coating: A Tool for Delivering Beneficial Microbes to Agricultural Crops. Frontiers in Plant Science 10: 1357

65. Singh R, Singh S, Parihar P, Mishra R K (2016) Reactive Oxygen Species) ROS (Beneficial Companions of Plants Developmental Processes. Frontiers in Plant Science 7:1299

66. Speakman J, Kruger W (1984) Control of Gaeumannomyces graminis var. tritici by a sterile, black mycelial fungus. Journal of Plant Diseases and Protection 91(5): 391-395.

67. Syamsia S, Idhan A, Firmansyah A P (2021) Combination on endophytic fungal as the plant growth-promoting fungi (PGPF) on cucumber (Cucumis sativus). Biodiversity Journal of Bio-Diversity 22(4):1194-1202.

68. Urja M and Meenu S (2010) Plant growth-promoting fungi: A review. Journal of Phytology 2(4): 33-44.

69. Zaidi N W, Dar M H, Singh S (2014) Trichoderma species as abiotic stress relievers in plants. In: Gupta V K, Schmoll M, Herrera-Estrella A, Upadhyay RS, Druzhinina I, Tuohy MG (eds) Biotechnology and Biology of Trichoderma. Elsevier, Amsterdam, pp 515-525.

70. Zhang F, Huo Y, Cobb A B (2018) Trichoderma biofertilizer links to altered soil chemistry, altered microbial communities, and improved grassland biomass. Frontiers in Microbiology 9:848

71. Zhang F, Meng X, Feng C et al (2016) Hydrolytic amino acids employed as a novel organic nitrogen source for the preparation of PGPF-containing bio-organic fertilizer for plant growth promotion and characterization of substance transformation during BOF production. PLoS One 11: 0149447

72. Zhang F, Zhu Z, Yang X, Ran W, Shen Q (2013) Trichoderma harzianum T-E5 significantly affects cucumber root exudates and fungal community in the cucumber rhizosphere. Applied Soil Ecology 72:41–48