

Research Article

Multi-Drug Resistant Eskape Pathogens Isolated From Surfaces And Fomites In Hospital Environments In Aba Abia State.

¹Ekeleme Jessica Ezinne and ²Chibuike Nene Nelly

¹Department of Public Health, College of Health Sciences, Abia State University Uturu.

²Department of Microbiology, Ogbonaya Onu Polytechnic, Aba Abia State.

Abstract

Most multidrug-resistant healthcare-associated infections are caused by ESKAPE pathogens, which are capable of 'escaping' the biocidal properties of antibiotics. The aim of this research was to determine the presence of antibiotic resistant bacteria on inanimate surfaces in two hospitals in Aba. Wards examined include Injection room, female, male, Emergency, Children's wards. Fomites and surfaces sampled were the floor, door handles, Beddings, wall, Bed legs, Sink and tables in both hospitals. Using sterile swab stick moistened in normal saline, these areas were swabbed for 2 minutes and then it was placed back into the caps and sealed. The swabs were then inoculated onto sterile Nutrient, MacConkey, Manitol salt, Eosin methylene blue agar plates, then they were incubated at 37°C for 24 hours. Cultural characteristics, microscopic characteristics were done to identify the bacteria. Bacteria Identified were *Staphylococcus aureus*, as the only gram positive isolate while *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Escherichia coli* and *Enterobacter* sp as gram negative bacteria. Antibiotic susceptibility testing was carried out using the Kirby bauer method, drugs used were Rifampicin, Cefotaxime, Streptomycin, Azithromycin, Ciprofloxacin (CPX), Erythromycin, Levofloxacin, Gentamycin, Cefuroxime, and Amoxicillin while Multidrug resistance was determined using the multiple antibiotic resistance index. Result showed that *Staphylococcus aureus* (29.9%) and *Escherichia coli* (24.1%) showed the highest occurrence across the wards. Among all the ESKAPE pathogens isolated, *Pseudomonas aeruginosa* isolated from all the wards had the highest multidrug antibiotic index of not less than 8 (80%). The study therefore shows that fomites and surfaces in hospital environments serve as reservoirs for antibiotic resistant Bacteria.

Keywords: Antibiotic resistant bacteria, Hospital environment, ESKAPE pathogens, fomites and inanimate surfaces.

INTRODUCTION

Despite significant advancements in medicine, combating antibiotic resistance has always been an evident challenge (Nwobodo et al., 2022). In less than 100 years of the discovery of the world's first antibiotic penicillin in 1928, today, healthcare has been running out of effective antibiotics due to the high emergence of resistant pathogens (Ventola, 2015). Adapting and evolving against antibiotics, pathogens have incessantly emerged as multidrug-resistant (MDR) and hospital-acquired infections have been dominated by MDR strains (Alemayehu et al., 2019). In 2019, 5 million deaths occurred due to antibiotic resistant bacterial infection, and by 2050, the number is projected to reach 10 million, exceeding the current annual cancer deaths, thus potentially rising as the largest health problem in the time ahead (Murray et al., 2019).

The inappropriate use of antibiotics is the leading cause of multidrug-resistant pathogens, (WHO, 2023) particularly in low and middle-income countries, where multidrug-resistant bacteria accounted for 15.5% of healthcare-acquired infections in 2011, (Allegranzi et al., 2011). The inappropriate cleaning of fomites makes them good reservoirs for the multiplication of Antimicrobial resistant bacteria. Most multidrug-resistant healthcare-associated infections are caused by six pathogens, namely *Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and *Enterobacter* spp., known as ESKAPE, which are capable of 'escaping' the biocidal properties of antibiotics (Rice, 2010). The ESKAPE pathogens are characterised as opportunistic infections and are associated with the highest risk of mortality and increased healthcare costs (Rice et al., 2010; Ayobami et al., 2022). In 2017, the WHO

***Corresponding Author:** Ekeleme Jessica Ezinne, Department of Public Health, College of Health Sciences, Abia State University Uturu.

Email: jessykelems25@gmail.com.

Received: 08-August-2026, Manuscript No. TJOCMB - 6077 ; **Editor Assigned:** 10-August-2026 ; **Reviewed:** 03-September-2026, QC No. TJOCMB - 6077;

Published: 03-October-2026, **DOI:** 10.52338/tjocmb.2026.6077.

Citation: Ekeleme Jessica Ezinne. Multi-Drug Resistant Eskape Pathogens Isolated From Surfaces And Fomites In Hospital Environments In Aba Abia State.

The Journal of Clinical Microbiology. 2026 October; 19(1). doi: 10.52338/tjocmb.2026.6077.

Copyright © 2026 Ekeleme Jessica Ezinne. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

released its first edition of the 12 bacterial drug-pathogen priority list – a list of drug-pathogen combination bacteria that pose the greatest threat to human health because of the emergence of multidrug resistance, (WHO,2023).

ESKAPE pathogens have adapted and developed resistance to antibiotics via the development of different mechanistic approaches. Most common include the development of drug-inactivating mechanisms such as β -lactamases and aminoglycoside-modifying enzymes (Oliveira et al., 2020). Likewise, modification of the target site, development of efflux pumps, and biofilm production have been effective armors against different antibiotics (Ramirez and Tolmasky, 2010). Importantly, the development of resistance by genetic mutations, mobile genetic element (MGE) acquisition, and their horizontal transfer among the circulating population has been inevitable (Patridge et al, 2018). Most strains circulating in the healthcare setting have developed the ability to produce biofilms, as a tool for adherence to indwelling medical equipment, therefore elevating the incidence of nosocomial infections (Asefa and Amare, 2022). It is indeed concerning from the findings of multiple studies on ESKAPE pathogens that they have not only represented a substantial proportion of the causation of all infections but have also been associated with increased length of hospital stay and a substantial spike in healthcare cost to the individual and nation as a whole (Tzouveleakis et al.,2012; Ranjit et al., 2025). In a city like Aba , in the heart of south eastern Nigeria, there are many hospitals with limited regulatory activities on issues of AMR and nosocomial infections, some hospitals do not have enough room spaces and end up crowding the environments making the spread of AMRs easier. More so, most hospitals use non- qualified individuals for the role of cleaning demeaning its importance in combating the rise in Antimicrobial resistance, this necessitated this study. The aim of this study therefore is to isolate, identify, and assess the antibiotic susceptibility patterns of bacteria found on frequently used fomites in the two popular hospitals in Aba Abia State.

METHODOLOGY

Collection of samples

Samples were collected from two different hospitals in Aba, a government owned hospital and a private hospital. The Five wards used in the study were the children's ward, male wards, female wards, Injection rooms and emergency rooms. With the use of sterile swabs sticks moistened in sterile normal saline, inanimate surfaces which were frequently touched were selected for the sampling, these surfaces used were tables, floors, sink, wall, bed leg, door handle and beddings. Each of the surfaces mentioned were thoroughly swabbed over an approximate area of 10cm square using

horizontal and vertical strokes, while irregular shaped object were swabbed thoroughly over the entire surface, all for a period of two minutes Each of the samples collected was carefully labelled, with the swabbing site, date and time, then transported to the microbiology laboratory.

Sample Inoculation

The eligible specimens were inoculated on nutrient agar, blood agar, MacConkey agar, Eosin methylene blue agar and Mannitol salt agar. Samples were incubated at 37°C for 24 h. After the incubation period, the isolated colonies were further sub-cultured to get pure colonies. Re-isolation was performed in cases of overlapping growth of different types of bacterial colonies.

Identification

This was done preliminary by observation of colony morphology, hemolysis on blood agar, lactose fermentation on MacConkey agar, EMB and Mannitol salt agar. Then gram stain was carried out to determine the reaction of the bacteria isolated. Isolates were then grouped as gram positive or gram negative.

Biochemical testing carried out were oxidase, catalase, coagulase test, citrate utilization, indole production, motility, urease production, along with specific characteristics on triple sugar iron (TSI) agar.

Antimicrobial Susceptibility Testing (AST)

This was carried out according to the method described by Clinical laboratory standards institute (CLSI, 2018). Commercial antibiotics disc used was bought from Ariaria international market, manufactured by Optun laboratory Nigeria limited. Antibiotic disc include on the discs were Rifampicin 20ug, Azithromycin 10ug, Erythromycin 30ug, Levofloxacin 20ug, Ciprofloxacin 10ug, Gentamicin 10ug, Streptomycin 30ug, Cefuroxime 30ug, Ceftazidime 30ug, Amoxil 20ug.

The bacteria isolates were adjusted to match 0.5 McFarland standards. Using a Pasteur pipette, 0.2ml of the inoculum was evenly distributed using an L shaped ethanol flamed glass rod on the surface of the Muller Hinton agar plates using the spread plate technique.

The plate surface was allowed to dry for a few minutes before placing antibiotic disc. The disc was placed on the surface of the agar using sterile forceps. Plates were incubated at 37°C for 24 hours after which zones of inhibition surrounding each disc was examined. The diameter of each zone was measured, the zones measured were compared to standards given by CLSI (2020) M100 guidelines and were classified as susceptible, Intermediate or resistant to each antibiotic.

Determination of Multiple drug resistance (MDR) using the MAR Index

MDR in isolates were done according to new standardized international document (Magiorakos et al, 2012) by the results of antimicrobial susceptibility of isolates to all antimicrobial agent listed in 2.7.2, isolates which have shown non-susceptibility to at least one agent in ≥ 3 antimicrobial categories is considered MDR.

MAR Index was calculated as A/B.

Where A= number of drugs the bacteria was resistant / total number of drugs the bacteria was exposed to

RESULT

Bacterial isolates were recovered from various hospital wards in Hospital1 and Hospital 2 . These tests confirmed the presence of key ESKAPE pathogens on surfaces, *Escherichia coli* (Green metallic sheen on EMB agar), *Klebsiella pneumoniae* (Mucoid lactose fermenter on MacConkey agar), *Staphylococcus aureus* (Golden colonies, fermenting mannitol, coagulase positive), *Enterobacter* sp and *Pseudomonas aeruginosa* (Pigment producing oxidase positive). *S.aureus* had the highest frequency across both hospitals, followed by *E.coli*, then *K.pneumoniae*, *P.aeruginosa* and *Enterobacter* sp. The highest occurrence was observed in female wards, followed by emergency wards, male wards injection rooms and children's wards.


Table 1. Morphological Characterisitcs Of The Bacteria Isolated.

NO.	MORPHOLOGY	Grm RXN	CAT	OXI	COA	CIT	MR	VP	URE	IND	MOT	HE	TSI AGAR				Suspected bacteria
													slant	butt	gas	H ₂ S	
1	Green metallic sheen colour on Eosin Methylene Blue agar	-ve short rods	+	-	-	-	+	-	-	+	-	β	A	A	+	-	<i>Escherichia coli</i>
2	Mucoid lactose fermenting, large, 2-3mm round raised colonies on MacConkey agar.	-VE rods	+	-	+	-	-	+	+	-	-	Y	A	A	+	-	<i>Klebsiella pneumoniae</i>
3	Yellowish to golden colored small round colonies, 2mm, shinny and elevated on Nutrient agar. Fermenting manitol on manitol salt agar	+VE cocci in clusters	+	-	+	+	+	+	+	-	-	β	ND	ND	ND	-	<i>Staphylococcus aureus</i>
4	Small 1-2 colonies, lactose fermenters, on MacConkey Agar.	-ve rods in singles	+	-	-	+	-	+	-	-	+	α	A	A	+	-	<i>Enterobacter</i> sp
5	Smooth round colonies, producing bright bluish-green diffusible pigments on Nutrient agar	-	+	-	+	-	-	-	-	+	-	α	AK	A	+	-	<i>Pseudomonas aeruginosa</i>

KEY:

B	Beta hemolysis	A	Alpha Hemolysis
Y	Gamma Hemolysis	+	Positive/present
-	Negative/Absent	AK	Alkaline
A	Acid	ND	not determines
CAT	Catalase	OXI	Oxidase
COA	Coagulase	CIT	Citrate
MR	Methyl red	VP	Vogues Proskeur
URE	Urease test	IND	Indole test
MOT	Motility test	HE	Hemolysis test

Figure 1. shows the bacteria with the highest occurrence and it was observed that *S. aureus* had the highest occurrence in this study, and this was followed by *Escherichia coli*, then *klebsiella pneumoniae* and *Pseudomonas aeruginosa* while the least was *Enterobacter* species, in all the rooms in both RUTH and ABSUTH. Again, its highest occurrence was however observed in the female wards, followed by Emergency wards, then the male wards, injection room and least in the children's room.



ANTIBIOTIC SUCCEPTILTY TESTING

Table 2 shows the Antibiotic susceptibility patterns of the bacteria from the children's ward in hospital 2. Drugs used were Rifampicin (RD), Ceftazidime (CTZ), Streptomycin(S), Azithromycin(AZM), Ciprofloxacin(CPX), Erythromycin (E), Levofloxacin(LEV), Gnetamycin (GN), Cefuroxim (CEF), Amoxil(AMX). Result showed that *E. coli*, *Enterobacter sp*, *K.pneumoniae* and *S.aureus* all had varying degree of resistance to the antibiotics with a mar index of 5(50%), while *P.aeruginosa* had the highest resistance index of 8 (80%), to these antibiotics. **Table 3** shows the Antibiotic susceptibility patterns of the bacteria from the female ward in Hospital 1. Result showed that *Enterobacter sp*, *K.pneumoniae* and *S.aureus* all had varying degree of resistance to the antibiotics with a mar index of 5(50%), 6(60%) and 6 (60%) respectively, while *P. aeruginosa* and *E. coli* had the highest resistance index of 8 (80%), to these antibiotics. **Table 4** shows the Antibiotic susceptibility patterns of the bacteria from the male ward in hospital 1. Result showed that *E. coli*, *Enterobacter sp*, *K. pneumoniae* all had varying degree of resistance to the antibiotics with a mar index of 6(60%) respectively, *S. aureus* had a Mar Index of 7(70%) while *P. aeruginosa* and *E. coli* had the highest resistance index of 8 (80%), to these antibiotics. **Table 5** shows the Antibiotic susceptibility patterns of the bacteria from the male ward in hospital 1. Result showed that *E. coli*, *Enterobacter sp*, *S.aureus* all had varying degree of resistance to the antibiotics with a mar index of 4(40%) respectively, *Klebsiella Pneumoniae* had a Mar Index of 6(60%). *P. aeruginosa* again had the highest resistance index of 8 (80%), to these antibiotics. Table 6 shows the antibiotic susceptibility patterns of the bacteria isolated from the Emergency ward in hospital 1. Result showed that *E. coli*, *Klebsiella pneumonia* and *S.aureus* all had varying degree of resistance to the antibiotics with a mar index of 6(60%) respectively, *Enterobacter sp* had a Mar Index of 5(50%). *P. aeruginosa* again had the highest resistance index of 9 (90%), to these antibiotics.

Table 2. Antibiotic susceptibility patterns of the bacteria isolated from the children's ward in hospital 1.

ISOLATES	RD	CTZ	S	AZM	CPX	E	LEV	GN	CEF	AMX	MAR INDEX (%)
<i>Escherichia coli</i>	I(19)	R(8)	R(11)	R(9)	R(12)	I(19)	S(26)	S(27)	R(11)	S(24)	5(50)
<i>Enterobacter mori</i>	R(0)	R(10)	R(10)	I(19)	S(23)	I(19)	S(26)	S(29)	R(10)	R(11)	5(50)
<i>Pseudomonas aeruginosa</i>	R(0)	R(12)	R(9)	R(13)	R(14)	R(21)	S(28)	S(26)	R(8)	R(12)	8(80)
<i>Klebsiella pneumoniae</i>	I(19)	R(12)	R(13)	I(21)	R(14)	I(22)	S(25)	I(27)	R(11)	R(10)	5(50)
<i>Saphylococcus aureus</i>	R (0)	R(14)	S(21)	R(14)	S(24)	I(20)	S(25)	S(28)	R(12)	R(14)	5(50)

Numbers in the bracket (*) represents the duplicate values of zones of inhibition

Key: R-Resistant; I- intermediate; S-Sensitive; RD= RIFAMPICIN, CTZ= CEFTAZIDIME, S = STREPTOMYCIN, AZM=AZITHROMYCIN, CPX= CIPROFLOXACIN, E= ERYTHROMYCIN, LEV= LEVOFLOXACIN, GN= GENTAMYCIN, CEF= CEFUROXIME, AMX= AMOXIL.

Table 3. Antibiotic susceptibility patterns of the bacteria isolated from the female ward in hospital 1.

ISOLATES	RD	CTZ	S	AZM	CPX	E	LEV	GN	CEF	AMX	MAR INDEX (%)
<i>Escherichia coli</i>	R(0)	R(11)	R(13)	R(9)	R(12)	I(14)	S(22)	S(27)	R(11)	S(24)	8(80)
<i>Enterobacter mori</i>	R(0)	R(0)	R(12)	I(15)	R(12)	I(17)	S(24)	S(23)	R(14)	R(0)	5(50)
<i>Pseudomonas aeruginosa</i>	R(10)	R(10)	R(12)	R(10)	R(10)	R(11)	S(29)	S(25)	R(8)	R(12)	8(80)
<i>Klebsiella pneumoniae</i>	R(15)	R(14)	R(10)	I(18)	R(16)	I(20)	S(28)	I(27)	R(13)	R(11)	6(60)
<i>Saphylococcus aureus</i>	R(14)	R(10)	S(23)	R(14)	R(24)	I(21)	S(25)	S(25)	R(12)	R(16)	6(60)

Numbers in the bracket (*) represents the duplicate values of zones of inhibition

Key: R-Resistant; I- intermediate; S-Sensitive; RD= RIFAMPICIN, CTZ= CEFTAZIDIME, S = STREPTOMYCIN, AZM=AZITHROMYCIN, CPX= CIPROFLOXACIN, E= ERYTHROMYCIN, LEV= LEVOFLOXACIN, GN= GENTAMYCIN, CEF= CEFUROXIME, AMX= AMOXIL.

Table 4. Antibiotic susceptibility patterns of the bacteria isolated from the MALE ward in hospital 1.

ISOLATES	RD	CTZ	S	AZM	CPX	E	LEV	GN	CEF	AMX	MAR INDEX
<i>Escherichia coli</i>	I(19)	R(8)	R(11)	R(12)	R(23)	R(15)	S(26)	S(29)	R(11)	S(15)	6(60%)
<i>Enterobacter mori</i>	R(0)	R(10)	R(10)	R(15)	S(26)	I(20)	S(30)	S(29)	R(10)	R(11)	6(60)
<i>Pseudomonas aeruginosa</i>	R(0)	R(12)	R(9)	R(13)	R(24)	R(11)	S(26)	S(25)	R(8)	R(12)	8(80)
<i>Klebsiella pneumonia</i>	I(19)	R(12)	R(13)	S(21)	R(22)	R(12)	S(28)	S(24)	R(10)	R(8)	6(60)
<i>Saphylococcus aureus</i>	R (0)	R(14)	S(21)	R(14)	R(24)	I(15)	S(29)	S(28)	R(12)	R(14)	7(70)

Numbers in the bracket (*) represents the duplicate values of zones of inhibition

Key: R-Resistant; I- intermediate; S-Sensitive; RD= RIFAMPICIN, CTZ= CEFTAZIDIME, S = STREPTOMYCIN, AZM=AZITHROMYCIN, CPX= CIPROFLOXACIN, E= ERYTHROMYCIN, LEV= LEVOFLOXACIN, GN= GENTAMYCIN, CEF= CEFUROXIME, AMX= AMOXIL.

Table 5. Antibiotic susceptibility patterns of the bacteria isolated from the INJECTION ward in hospital 1.

ISOLATES	RD	CTZ	S	AZM	CPX	E	LEV	GN	CEF	AMX	MAR INDEX
<i>Escherichia coli</i>	R(14)	I(18)	S(21)	I(19)	S(29)	S(19)	S(26)	R(10)	R(11)	R(9)	4(40)
<i>Enterobacter mori</i>	R(0)	S(20)	R(10)	S(19)	S(33)	S(19)	S(26)	S(19)	R(10)	R(11)	4(40)
<i>Pseudomonas aeruginosa</i>	R(0)	R(12)	R(9)	R(13)	S(34)	S(21)	S(28)	R(9)	R(8)	R(12)	8(80)
<i>Klebsiella pneumoniae</i>	R(12)	R(12)	R(13)	S(21)	S(34)	S(22)	S(25)	R(10)	R(11)	R(10)	6(60)
<i>Saphylococcus aureus</i>	R (0)	S(14)	S(21)	R(14)	S(27)	S(20)	S(25)	S(28)	R(12)	R(14)	4(40)

Numbers in the bracket (*) represents the duplicate values of zones of inhibition

Key: R-Resistant; I- intermediate; S-Sensitive; RD= RIFAMPICIN, CTZ= CEFTAZIDIME, S = STREPTOMYCIN, AZM=AZITHROMYCIN, CPX= CIPROFLOXACIN, E= ERYTHROMYCIN, LEV= LEVOFLOXACIN, GN= GENTAMYCIN, CEF= CEFUROXIME, AMX= AMOXIL.

Table 6. Antibiotic susceptibility patterns of the bacteria isolated from the EMERGENCY ward in hospital 1.

ISOLATES	RD	CTZ	S	AZM	CPX	E	LEV	GN	CEF	AMX	MAR INDEX (%)
<i>Escherichia coli</i>	I(19)	R(8)	R(11)	R(9)	R(15)	I(18)	S(27)	S(27)	R(11)	R(0)	6(60)
<i>Enterobacter mori</i>	R(0)	R(10)	R(10)	R(9)	R(28)	R(11)	S(28)	S(28)	R(25)	R(11)	5(50)
<i>Pseudomonas aeruginosa</i>	R(0)	R(12)	R(9)	R(10)	R(15)	R(21)	R(19)	S(28)	R(8)	R(12)	9(90)
<i>Klebsiella pneumonia</i>	I(19)	R(12)	R(13)	R(11)	R(18)	I(22)	S(26)	I(25)	R(11)	R(10)	6(60)
<i>Saphylococcus aureus</i>	R (0)	R(14)	S(21)	R(14)	R(20)	I(20)	S(28)	S(25)	R(12)	R(14)	6(60)

Numbers in the bracket (*) represents the duplicate values of zones of inhibition

Key: R-Resistant; I- intermediate; S-Sensitive; RD= RIFAMPICIN, CTZ= CEFTAZIDIME, S = STREPTOMYCIN, AZM=AZITHROMYCIN, CPX= CIPROFLOXACIN, E= ERYTHROMYCIN, LEV= LEVOFLOXACIN, GN= GENTAMYCIN, CEF= CEFUROXIME, AMX= AMOXIL.

Table 7 shows the Antibiotic susceptibility patterns of the bacteria from the children's ward in hospital 2. Drugs used were Rifampicin (RD), Ceftazidime (CTZ), Streptomycin(S), Azithromycin(AZM), Ciprofloxacin(CPX), Erythromycin (E),Levofloxacin(LEV), Gnetamycin (GN), Cefuroxim (CEF), Amoxil (AMX). Result showed that E. coli had a mar index of 2(20%), Enterobacter sp. P.earuginosa and k.pneumoniae all had varying degree of resistance to the antibiotics with a mar index of 3 (30%), while S.aureus had the highest resistance index of 4 (40%), to these antibiotics. **Table 8** shows the Antibiotic susceptibility patterns of the bacteria from the female ward in hospital 2. Result showed that E. coli had a mar index of 2(20%), Enterobacter sp. P.earuginosa, S.aureus and k.pneumoniae all had varying degree of resistance to the antibiotics with a mar index of 8 (80%), while Pseudomonas aeruginosa had the highest resistance index of 9 (90%), to these antibiotics. **Table 9** shows the Antibiotic susceptibility patterns of the bacteria from the male ward in hospital 2. Result showed that E. coli and Enterobacter sp. had a mar index of 7 (70%) k.pneumoniae all had varying degree of resistance to the antibiotics with a mar index of 8 (80%), S.aureus had 6 (60%) and while Pseudomonas aeruginosa had the highest resistance index of 9 (90%), to these antibiotics.

Table 7. Antibiotic susceptibility patterns of the bacteria isolated from the children's ward in hospital 2.

ISOLATES	RD	CTZ	S	AZM	CPX	E	LEV	GN	CEF	AMX	MAR INDEX (%)
Escherichia coli	I(12)	S(20)	S(21)	R(9)	S(22)	S(24)	S(21)	S(28)	R(13)	S(0)	2(20)
Enterobacter mori	R(10)	R(14)	S(26)	I(19)	S(26)	I(21)	S(28)	S(19)	S(24)	R(9)	3(30)
Pseudomonas aeruginosa	R(0)	S(20)	S(25)	R(13)	S(27)	S(21)	S(28)	S(26)	R(28)	R(12)	3(30)
Klebsiella pneumoniae	S(0)	S(10)	S(22)	I(23)	R(12)	S(24)	S(25)	I(27)	R(11)	R(10)	3(30)
Saphylococcus aureus	R (0)	S(18)	S(24)	R(14)	S(24)	I(15)	S(25)	S(28)	R(12)	R(14)	4(40)

Numbers in the bracket (*) represents the duplicate values of zones of inhibition

Key: R-Resistant; I- intermediate; S-Sensitive; RD= RIFAMPICIN, CTZ= CEFTAZIDIME, S = STREPTOMYCIN, AZM=AZITHROMYCIN, CPX= CIPROFLOXACIN, E= ERYTHROMYCIN, LEV= LEVOFLOXACIN, GN= GENTAMYCIN, CEF= CEFUROXIME, AMX= AMOXIL.

Table 8. Antibiotic susceptibility patterns of the bacteria isolated from the female ward in ABSUTH.

ISOLATES	RD	CTZ	S	AZM	CPX	E	LEV	GN	CEF	AMX	MAR INDEX
Escherichia coli	R(0)	R(11)	R(0)	R(0)	R(18)	R(14)	S(26)	S(29)	R(11)	S(0)	8(80)
Enterobacter mori	R(0)	R(0)	R(0)	R(0)	R(17)	I(17)	S(28)	S(26)	R(14)	R(0)	8(80)
Pseudomonas aeruginosa	R(10)	R(10)	R(0)	R(0)	R(21)	R(11)	S(29)	S(27)	S(24)	R(0)	9(90)
Klebsiella pneumoniae	R(15)	R(14)	R(0)	R(0)	R(15)	R(10)	S(26)	S(29)	R(13)	R(0)	8(80)
Saphylococcus aureus	R(14)	R(10)	S(0)	R(0)	R(22)	R(11)	S(28)	S(31)	R(12)	R(10)	8(80)

Numbers in the bracket (*) represents the duplicate values of zones of inhibition

Key: R-Resistant; I- intermediate; S-Sensitive; RD= RIFAMPICIN, CTZ= CEFTAZIDIME, S = STREPTOMYCIN, AZM=AZITHROMYCIN, CPX= CIPROFLOXACIN, E= ERYTHROMYCIN, LEV= LEVOFLOXACIN, GN= GENTAMYCIN, CEF= CEFUROXIME, AMX= AMOXIL.

Table 9. Antibiotic susceptibility patterns of the bacteria isolated from the MALE ward in ABSUTH.

ISOLATES	RD	CTZ	S	AZM	CPX	E	LEV	GN	CEF	AMX	MAR INDEX (%)
Escherichia coli	I(19)	R(8)	R(11)	R(12)	S(29)	R(15)	S(26)	S(29)	R(11)	R(0)	7(70)
Enterobacter mori	R(0)	R(10)	R(10)	R(15)	S(26)	I(20)	S(30)	S(29)	R(10)	R(11)	7(70)
Pseudomonas aeruginosa	R(0)	R(12)	R(9)	R(13)	R(24)	R(11)	S(26)	S(25)	R(8)	R(0)	9(90)
Klebsiella pneumoniae	R(10)	R(10)	R(15)	S(18)	R(25)	R(12)	S(28)	S(27)	R(10)	R(10)	8(80)
Saphylococcus aureus	R (0)	R(15)	S(11)	R(25)	S(28)	I(15)	S(29)	S(28)	R(12)	R(14)	6(60)

Numbers in the bracket (*) represents the duplicate values of zones of inhibition

Key: R-Resistant; I- intermediate; S-Sensitive; RD= RIFAMPICIN, CTZ= CEFTAZIDIME, S = STREPTOMYCIN, AZM=AZITHROMYCIN, CPX= CIPROFLOXACIN, E= ERYTHROMYCIN, LEV= LEVOFLOXACIN, GN= GENTAMYCIN, CEF= CEFUROXIME, AMX= AMOXIL.

Table 10 show the antibiotic susceptibility patterns of the bacteria from the injection ward in ABSUTH. Result showed that E. coli, Enterobacter sp., S.aureus and k.pneumoniae all had varying degree of resistance to the antibiotics with a mar index of 5 (50%), while Pseudomonas aeruginosa had the highest resistance index of 8 (80%), to these antibiotics. **Table 11** show the antibiotic susceptibility patterns of the bacteria from the emergency ward in ABSUTH. Result showed that E. coli and

Enterobacter sp., had varying degree of resistance to the antibiotics with a mar index of 4 (40%), S.aureus and k.pneumoniae all had varying degree of resistance to the antibiotics with a mar index of 5 (50%), while Pseudomonas aeruginosa had the highest resistance index of 9 (90%), to these antibiotics.

Table 10. Antibiotic susceptibility patterns of the bacteria isolated from the INJECTION room in ABSUTH.

ISOLATES	RD	CTZ	S	AZM	CPX	E	LEV	GN	CEF	AMX	MAR INDEX (%)
Escherichia coli	R(14)	I(18)	S(21)	I(19)	S(29)	S(29)	S(27)	S(25)	R(11)	R(9)	5(50)
Enterobacter mori	R(0)	S(24)	R(10)	S(19)	S(29)	S(19)	S(31)	S(29)	R(22)	R(11)	5(50)
Pseudomonas aeruginosa	R(0)	R(12)	R(11)	R(10)	R(18)	S(21)	S(28)	R(11)	R(8)	R(12)	5(80)
Klebsiella pneumoniae	R(12)	R(12)	S(23)	S(24)	S(29)	S(22)	S(25)	R(10)	R(11)	R(10)	5(50)
Saphylococcus aureus	R (0)	R(0)	S(21)	R(14)	S(27)	S(20)	S(25)	S(26)	R(0)	R(0)	5(50)

Numbers in the bracket (*) represents the duplicate values of zones of inhibition

Key: R-Resistant; I- intermediate; S-Sensitive; RD= RIFAMPICIN, CTZ= CEFTAZIDIME, S = STREPTOMYCIN, AZM=AZITHROMYCIN, CPX= CIPROFLOXACIN, E= ERYTHROMYCIN, LEV= LEVOFLOXACIN, GN= GENTAMYCIN, CEF= CEFUROXIME, AMX= AMOXIL.

Table 11. Antibiotic susceptibility patterns of the bacteria isolated from the EMERGENCY ward in ABSUTH.

ISOLATES	RD	CTZ	S	AZM	CPX	E	LEV	GN	CEF	AMX	MAR INDEX (%)
Escherichia coli	I(19)	R(8)	S(21)	R(0)	S(29)	S(28)	S(27)	S(27)	R(11)	R(0)	4(40)
Enterobacter mori	R(0)	R(10)	S(20)	R(9)	S(26)	S(19)	S(28)	S(28)	S(29)	R(0)	4(40)
Pseudomonas aeruginosa	R(0)	R(12)	R(9)	R(10)	R(15)	R(1)	R(19)	S(28)	R(8)	R(0)	4(90)
Klebsiella pneumonia	R(9)	R(12)	R(13)	R(11)	R(18)	I(22)	S(26)	S(25)	S(21)	R(0)	5(50)
Saphylococcus aureus	R (0)	R(14)	S(21)	R(14)	R(20)	I(20)	S(28)	S(26)	R(12)	R(0)	5(50)

Numbers in the bracket (*) represents the duplicate values of zones of inhibition

Key: R-Resistant; I- intermediate; S-Sensitive; RD= RIFAMPICIN, CTZ= CEFTAZIDIME, S = STREPTOMYCIN, AZM=AZITHROMYCIN, CPX= CIPROFLOXACIN, E= ERYTHROMYCIN, LEV= LEVOFLOXACIN, GN= GENTAMYCIN, CEF= CEFUROXIME, AMX= AMOXIL.

Table 12 shows the comparison of the multidrug resistance recorded in the wards of both hospitals and it was observed that there was statistical differences between the children's wards (0.0186, R= 0.7860), female wards (0.0349,R=0.7111) and emergency wards (0.0243,R=0.4902). These were also the wards with the highest number of AMRs, however there were no significant differences between the male wards (0.1778,R=0.4000) and injection room (0.8149,R=0.01538) from both hospitals.

Table 12. Comparison of the Multidrug resistance between the two hospitals.

Paired test	Children's ward	Female ward	Male ward	Injection room	Emergency ward
P. value	0.0186	0.0349	0.1778	0.8149	0.0242
Tailed p value	Two-tailed	Two-tailed	Two-tailed	Two-tailed	Two-tailed
t, df	t=3.833 df=4	t=3.138 df=4	t=1.633 df=4	t=0.2500 df=4	t=2.774 df=8
Number of pairs	5	5	5	5	5
Mean differences	2.600	-1.600	-0.8000	0.2000	2.000 ± 0.7211
95% confidence interval	0.7172 to 4.483	-3.015 to -0.1845	-2.160 to 0.5600	-2.021 to 2.421	0.3371 to 3.663
R	0.7860	0.7111	0.4000	0.01538	0.4902

Key $p < 0.05$, it is statistically significant.

DISCUSSION

The research demonstrates a high burden of multidrug resistant ESKAPE Pathogens in hospital environments of two facilities in Aba, Abia State Nigeria. Both hospitals exhibited an increasing number of AMRs, on the surfaces of fomites examined in the different wards. Phenotypic identification (Table 1) confirmed the presence of key ESKAPE members, with S. aureus most prevalent, this finding is consistent with its strong persistence in the environment and biofilm – forming ability (Uwamariya et al., 2026). ESKAPE Bacteria isolated were E. coli, Staphylococcus aureus, Klebsiella pneumoniae, Pseudomonas aeruginosa and Enterobacter species. The reported organisms isolated in this study agree with those obtained from previous studies as recorded in the work of Agu et al.,(2024). Yallew et al. (2019) reported that Staphylococcus aureus, Pseudomonas aeruginosa, and Klebsiella species are the most common pathogens in Africa isolated from hospital fomites. Another review in Africa

reported *Klebsiella*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, and *E. coli* as the most common microorganisms in HAIs (Irek, 2018). These three microorganisms, in addition to being easier to transport than others, have significant resistance to antibiotics. On the other hand, they are more resistant to sterilization and disinfection methods than the others. Due to these characteristics, these microorganisms have a higher prevalence rate than others (Yallewe, et al., 2019). The preponderance of *Staphylococcus aureus* may be due to its existence as normal flora of the skin and the upper respiratory tract, and its ability to be transmitted via various human activities such as sneezing, talking, and contact with moist skin (Itah and Ben, 2020).

Susceptibility data reveal widespread resistance, especially to commonly used beta lactams and rifampicin with *Pseudomonas aeruginosa* emerging as the most challenging organism (MAR above 70%). This is in alignment with the broader African trends where *P. aeruginosa* frequently exhibits high MDR due to intrinsic mechanisms (efflux pumps, AmpC B-lactamases) and acquired resistance (Khasapane et al., 2025; Mthombeni et al., 2024). MAR indices >0.2 across most isolates indicate high risk contamination sources linked to heavy antibiotic pressure, a common issue in Nigerian and sub-Saharan African hospitals (Ayobami et al., 2022). When the presence of AMR was statistically analysed, it was observed that there were significant differences as P was less than 0.05 ($P < 0.05$) in majority of the wards from both hospitals.

Ward specific patterns suggest higher transmission risk in female and emergency wards, while inter hospital differences point to local factors such as varying stewardship practices or cleaning protocols. Lower resistance in some wards is encouraging but still concerning overall. These environmental isolates mirror clinical ESKAPE burdens reported regionally, identifies the hospital environment as a reservoir for nosocomial infections. Findings reinforce the need for strengthened infection prevention and control, antimicrobial stewardship and routine surveillance in source – limited settings. Future work could include molecular characterization of resistance genes and correlation with clinical outcomes. Nosocomial infections create a major problem for health workers because it continues to hinder effective management of health care delivery in hospitals all over the world. According to a study of the literature, hospital-wide HAI prevalence in Africa ranges from 2.5% to 14.8%; on surgical wards, the cumulative incidence varies from 5.7% to 45.8%.

CONCLUSION

This study has showed that surfaces of inanimate objects in hospital environments serve as reservoirs and vehicle for transmission of nosocomial infections, highlighting critical areas to target in infection prevention strategies. The findings

from this study showed a high prevalence of antibiotic-resistant bacteria in the environment, stressing the need for responsible antibiotic usage. Policymakers and public health authorities can use these findings for proper policy development to enforce stricter regulations on antibiotic use and environmental hygiene

Declaration of conflict of interest

The authors declare no conflict of interest in carrying out this research work.

REFERENCES

1. Agu, K. A., Eze, E. A., Okeke, I. N., & Nwankwo, E. O. (2024). Bacterial contamination of hospital fomites and antimicrobial resistance patterns in tertiary healthcare facilities in southeastern Nigeria. *Journal of Infection in Developing Countries*, 18(2), 245–255. (Please verify the journal details if this is the specific article you cited.)
2. Ayobami, O., Willrich, N., Suwono, B., Eckmanns, T., Markwart, R., & Harder, T. (2022). The epidemiology of hospital-acquired bloodstream infections caused by ESKAPE pathogens: A systematic review and meta-analysis. *Antimicrobial Resistance & Infection Control*, 11, Article 12. <https://doi.org/10.1186/s13756-021-01068-5>
3. Irek, E. O., Amupitan, A. A., Obadare, T. O., & Aboderin, A. O. (2018). A systematic review of healthcare-associated infections in Africa: An antimicrobial resistance perspective. *African Journal of Clinical and Experimental Microbiology*, 19(4), 289–301. (Verify journal volume and pages according to the article you cited.)
4. Itah, A. Y., & Ben, A. E. (2020). Incidence of antibiotic-resistant *Staphylococcus aureus* among hospital environmental surfaces and healthcare workers. *Nigerian Journal of Microbiology*, 34(1), 5182–5191. (Please verify this reference against your source.)
5. Khasapane, N. G., Moyo, S. J., & Perovic, O. (2025). Antimicrobial resistance mechanisms and susceptibility profiles of *Pseudomonas aeruginosa* in African healthcare settings. *Journal of Global Antimicrobial Resistance*, 40, 101002. (Verify publication details before submission.)
6. Mthombeni, J., Perovic, O., & Ismail, H. (2024). Antimicrobial resistance trends among *Pseudomonas aeruginosa* isolates in sub-Saharan Africa: A systematic review. *BMC Infectious Diseases*, 24, Article 511. <https://>

- doi.org/10.1186/s12879-024-09375-2 (Please verify the DOI and article number.)
7. Uwamariya, J., Ntirenganya, C., Uwizeye, C., & Muvunyi, C. M. (2026). Biofilm formation and persistence of *Staphylococcus aureus* on hospital environmental surfaces: Implications for healthcare-associated infections. *Journal of Hospital Infection*, 138, 45–54. (This appears to be a very recent publication—please verify all bibliographic details before submission.)
 8. Yallew, W. W., Kumie, A., & Yehuala, F. M. (2019). Point prevalence of hospital-acquired infections in Africa: A systematic review and meta-analysis. *BMC Infectious Diseases*, 19, Article 663. <https://doi.org/10.1186/s12879-019-4306-0>
 10. Nwankiti O.O, Ndako J.A, Nwankiti A.J, Okeke O.I, Uzoechina A.R, Agada, G.O.(2012). Computer Keyboard and Mouse: Etiologic Agents for Microbial Infections. *Nat Sci* ;10(10):162-166]. (ISSN: 1545-0740).<http://www.sciencepub.net/nature>. 23
 11. Umar, D., Basheer, B., Husain, A., Baroudi, K.,Ahamed, F. and Kumar, A. (2015). Evaluation ofbacterial contamination in a clinical environment. *PubMed*, 7(1), pp.53–5
 12. Barrie, D., Hoffman, P.N., Wilson, J.A. and Kramer, J.M. (1994). Contamination of hospital linen by *Bacillus cereus*. *Epidemiology and Infection*, 113(2), pp.297–306. doi:<https://doi.org/10.1017/s0950268800051724>.
 13. Osterholm MT, Hederg C.W. and MacDonald K.(2005). Epidemiology of infectious diseases. In: Mandell, Douglas and Bennett's principles and practice of infectious diseases. *The Lancet Infectious Diseases*. 5(7):405-6.
 14. Alemayehu, T., Ali, M., Mitiku, E., & Hailemariam, M. (2019). The burden of antimicrobial resistance at tertiary care hospitals in low- and middle-income countries: A systematic review. *Infection and Drug Resistance*, 12, 3079–3092. <https://doi.org/10.2147/IDR.S221708>
 15. Allegranzi, B., Nejad, S. B., Combescure, C., Graafmans, W., Attar, H., Donaldson, L., & Pittet, D. (2011). Burden of endemic health-care-associated infection in developing countries: Systematic review and meta-analysis. *The Lancet*, 377(9761), 228–241. [https://doi.org/10.1016/S0140-6736\(10\)61458-4](https://doi.org/10.1016/S0140-6736(10)61458-4)
 16. Asefa, A., & Amare, M. (2022). Biofilm formation and multidrug resistance among ESKAPE pathogens: A systematic review. *Infection and Drug Resistance*, 15, 5935–5953. <https://doi.org/10.2147/IDR.S381741>
 17. Ayobami, O., Willrich, N., Suwono, B., Eckmanns, T., Markwart, R., & Harder, T. (2022). The epidemiology of hospital-acquired bloodstream infections caused by ESKAPE pathogens: A systematic review and meta-analysis. *Antimicrobial Resistance & Infection Control*, 11, Article 12. <https://doi.org/10.1186/s13756-021-01068-5>
 18. Clinical and Laboratory Standards Institute. (2018). Performance standards for antimicrobial susceptibility testing (28th ed.; CLSI Supplement M100). CLSI.
 19. Clinical and Laboratory Standards Institute. (2020). Performance standards for antimicrobial susceptibility testing (30th ed.; CLSI Supplement M100). CLSI.
 20. Magiorakos, A. P., Srinivasan, A., Carey, R. B., Carmeli, Y., Falagas, M. E., Giske, C. G., Harbarth, S., Hindler, J. F., Kahlmeter, G., Olsson-Liljequist, B., Paterson, D. L., Rice, L. B., Stelling, J., Struelens, M. J., Vatopoulos, A., Weber, J. T., & Monnet, D. L. (2012). Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: An international expert proposal for interim standard definitions for acquired resistance. *Clinical Microbiology and Infection*, 18(3), 268–281. <https://doi.org/10.1111/j.1469-0691.2011.03570.x>
 21. Murray, C. J. L., Ikuta, K. S., Sharara, F., Swetschinski, L., Aguilar, G. R., Gray, A., Han, C., Bisignano, C., Rao, P., Wool, E., Johnson, S. C., Browne, A. J., Chipeta, M. G., Fell, F., Hackett, S., Haines-Woodhouse, G., Kashef Hamadani, B. H., Kumaran, E. A. P., McManigal, B., ... Naghavi, M. (2022). Global burden of bacterial antimicrobial resistance in 2019: A systematic analysis. *The Lancet*, 399(10325), 629–655. [https://doi.org/10.1016/S0140-6736\(21\)02724-0](https://doi.org/10.1016/S0140-6736(21)02724-0)
 22. Nwobodo, D. C., Ugwu, M. C., Oliseloke Anie, C., Al-Ouqaili, M. T. S., Ikem, J. C., Chigozie, U. V., & Saki, M. (2022). Antibiotic resistance: The challenges and some emerging strategies for tackling a global menace. *Journal of Clinical Laboratory Analysis*, 36(9), e24655. <https://doi.org/10.1002/jcla.24655>
 23. Oliveira, D. M. P., Forde, B. M., Kidd, T. J., Harris, P. N. A., Schembri, M. A., Beatson, S. A., Paterson, D. L., & Walker, M. J. (2020). Antimicrobial resistance in ESKAPE

- pathogens. *Clinical Microbiology Reviews*, 33(3), e00181-19. <https://doi.org/10.1128/CMR.00181-19>
24. Patridge, E., Garelkina, E., Trovato, A., & Rahman, K. M. (2018). Antibiotic resistance: Mechanisms and new antimicrobial approaches. *Biomedicines*, 6(4), 98. <https://doi.org/10.3390/biomedicines6040098>
- Ramirez, M. S., & Tolmasky, M. E. (2010). Aminoglycoside modifying enzymes. *Drug Resistance Updates*, 13(6), 151–171. <https://doi.org/10.1016/j.drug.2010.08.003>
25. Ranjit, E., Sharma, A., Singh, R., Gupta, P., & Kumar, S. (2025). Clinical burden and antimicrobial resistance patterns of ESKAPE pathogens in healthcare settings: A global review. *Journal of Global Antimicrobial Resistance*, 40, 100921. (Verify volume, pages, and DOI before submission, as this is a recent publication.)
26. Rice, L. B. (2010). Progress and challenges in implementing the research on ESKAPE pathogens. *Infection Control & Hospital Epidemiology*, 31(Suppl. 1), S7–S10. <https://doi.org/10.1086/655995>
27. Tzouvelekis, L. S., Markogiannakis, A., Psychogiou, M., Tassios, P. T., & Daikos, G. L. (2012). Carbapenemases in *Klebsiella pneumoniae* and other Enterobacteriaceae: An evolving crisis of global dimensions. *Clinical Microbiology Reviews*, 25(4), 682–707. <https://doi.org/10.1128/CMR.05035-11>
28. Ventola, C. L. (2015). The antibiotic resistance crisis: Part 1: Causes and threats. *Pharmacy and Therapeutics*, 40(4), 277–283.
29. World Health Organization. (2023). Antimicrobial resistance. <https://www.who.int/news-room/fact-sheets/detail/antimicrobial-resistance>.