

Effect of garlic aqueous extract on liver protection against liver toxicity brought on by ciprofibrate in male albino rats

Azab Elsayed Azab

*Corresponding author

Azab Elsayed Azab, Department of Physiology, Faculty of Medicine, Sabratha University, Libya.

Received Date : July 15, 2022

Accepted Date : July 16, 2022

Published Date : Aug 16, 2022

Abstract

Background : The danger of liver growth is increased by the liver's everyday exposure to too many hazardous compounds, either from the environment or from a variety of other sources like preservatives or medications. One of the chemical families of peroxisome proliferators that stimulates hepatic cells and hepatic cell proliferation is ciprofibrate. uncontrollably divides, resulting in liver expansion. Drugs and other xenobiotics, such as the long-term hypolipidemic treatment that causes cancer in rats, can cause the liver cell to divide. However, this has not yet been proven in people. The current study's objectives are to assess the possible protective benefits of garlic administration against the biochemical and histological changes that ciprofibrate-induced liver damage in male rats.

Materials and Procedures : In the current investigation, 6 male rats were divided into 8 groups: Control group, Oil, Garlic, Ciprofibrate 50 and 100 mg/kg body weight, Cipro 50 mg/kg body weight plus garlic, Cipro 100 mg/kg body weight plus garlic. For 21 days, the rats received daily oral gavage treatments. The animals were killed on the final day of the experiment, and then blood samples and liver tissue were taken. Serum Alanine Aminotransferase (ALT), Aspartate Aminotransferase (AST), Alkaline Phosphates (ALP), and serum total and direct bilirubin concentrations were measured to assess liver function. For all groups, a histological analysis of the liver tissues was done.

Conclusion : Treatment with ciprofibrate led to significant histopathological and liver function test elevations, and garlic aqueous extract was able to shield male rats' livers from these side effects. Therefore, it should be recommended to patients who are on ciprofibrate treatment to take garlic aqueous extract.

Discussion

The liver is a key organ for detoxification and a main location for intensive metabolism in general, making it susceptible to a variety of illnesses. as a result of being exposed to the poisons . In recent decades, liver function tests have emerged as the most crucial tools for evaluating the liver's resistance to damage. The severity of the liver injury is often assessed by liver performance indices such ALT, AST, and ALP . In hepatotoxicity investigations brought on by chemicals, serum aminotransferase activities are known as toxicity markers. An increase in these enzymes' activities is referred to as the early detection of toxic hepatitis. a considerable amount of such peroxisome proliferators. It is important to note that free radicals injure healthy cells and contribute significantly to the occurrence of cancer, but garlic, which is a potent antioxidant, lessens the harm done by these agents. In this investigation, garlic extract dramatically decreased serum liver functions and had a beneficial impact on cell proliferation. Proliferation, a process that leads to an increase in the number of cells, is balanced between cell division and cell loss through cell death or differentiation. It reduced the rats treated with ciprofibrate's liver damage. Garlic regulates how cells use oxygen, and it appears that the protective effects of garlic extract entail the preservation of antioxidant capacity in preventing oxidative stress and safeguarding hepatic tissue.

Conclusion

The current study looked at the protective benefits of garlic against liver damage brought on by peroxisome proliferators (ciprofibrate). The prospective effects of ciprofibrate on liver enzymes and the histological consequences are accurately analysed in this study. Through examination of biochemical markers and histological alterations, the study produced valuable data. Peroxisome proliferators have been the subject of prior research, and the current study largely supported those findings. Garlic was also utilised in this experiment to lessen the drug's toxicological effects. In rats, liver damage brought on by hepatotoxicity was inhibited by garlic.

References

1. Banerjee, R. R. et al. (2004) Regulation of fasted blood glucose by resistin', Science. American Association for the Advancement of Science, 303(5661), pp. 1195-1198.

2. Hyder, M. A., Hasan, M. and Mohieldein, A. H. (2013) 'Comparative levels of ALT, AST, ALP and GGT in liver associated diseases', *European journal of experimental biology*, 3(2), pp. 280–284.
3. Vanden Heuvel, J. P. et al. (2006) 'Differential activation of nuclear receptors by per fluorinated fatty acid analogs and natural fatty acids: a comparison of human, mouse, and rat peroxisome proliferators-activated receptor- α , - β , and - γ , liver X receptor- β , and retinoid X receptor- α ', *Toxicological Sciences*. Oxford University Press, 92(2), pp. 476–489.
4. Darnell Jr, J. E. (2002) 'Transcription factors as targets for cancer therapy', *Nature Reviews Cancer*. Nature Publishing Group, 2(10), p. 740.
5. Abushofa, F. A. A. (2014) 'Studies on the role of peroxisome proliferators: in liver growth and neurodegenerative disorders'. University of Nottingham.
6. Vanhove, G. F. et al. (1993) 'The CoA esters of 2-methyl-branched chain fatty acids and of the bile acid intermediates di-and trihydroxy-coprostanic acids are oxidized by one single peroxisomal branched chain acyl-CoA oxidase in human liver and kidney.', *Journal of Biological Chemistry*. ASBMB, 268(14), pp. 10335–10344.
7. Kliewer, S. A. et al. (1997) 'Fatty acids and eicosanoids regulate gene expression through direct interactions with peroxisome proliferator-activated receptors α and γ ', *Proceedings of the National Academy of Sciences*. National Acad Sciences, 94(9), pp. 4318–4323.
8. Mannaerts, G. P. and Van Veldhoven, P. P. (1993) 'Metabolic pathways in mammalian peroxisomes', *Biochimie*. Elsevier, 75(3–4), pp. 147–158.
9. Rao, M. S. and Subbarao, V. (1997) 'Effect of dexamethasone on ciprofibrate-induced cell proliferation and peroxisome proliferation', *Toxicological Sciences*. Oxford University Press, 35(1), pp. 78–83.
10. Al Kholaiifi, A. et al. (2008) 'Species-specific kinetics and zonation of hepatic DNA synthesis induced by ligands of PPAR α ', *Toxicological sciences*. Oxford University Press, 104(1), pp. 74–85.
11. Kawanishi, S. et al. (2002) 'The role of metals in site-specific DNA damage with reference to carcinogenesis', *Free Radical Biology and Medicine*. Elsevier, 32(9), pp. 822–832.
12. Tripathi, K. (2009) 'A review-Garlic, the spice of life-(Part-I)', *Asian J Res Chem*, 2(1), pp. 8–13.
13. Shaarawy, S. M. et al. (2009) 'Protective effects of garlic and silymarin on NDEA-induced rats hepatotoxicity', *International journal of biological sciences*. Ivyspring International Publisher, 5(6), p. 549.
14. Agarwal, K. C. (1996) 'Therapeutic actions of garlic constituents', *Medicinal research reviews*. Wiley Online Library, 16(1), pp. 111–124.
15. Aruoma, O. I. et al. (1992) 'Antioxidant and prooxidant properties of active rosemary constituents: carnosol and carnosic acid', *Xenobiotica*. Taylor & Hepatoprotective Effect of Garlic Aqueous Extract against Hepatotoxicity Induced By Ciprofibrate in Male Albino Rats Francis, 22(2), pp. 257–268.
16. Kita, T. et al. (1987) 'Probucol prevents the progression of atherosclerosis in Watanabe heritable hyperlipidemic rabbit, an animal model for familial hypercholesterolemia', *Proceedings of the National Academy of Sciences*. National Acad Sciences, 84(16), pp. 5928–5931.
17. Forman, B. M., Chen, J. and Evans, R. M. (1997) 'Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors α and δ ', *Proceedings of the National Academy of Sciences*. National Acad Sciences, 94(9), pp. 4312–4317.
18. Bordia, A. (1981) 'Effect of garlic on blood lipids in patients with coronary heart disease', *The American Journal of Clinical Nutrition*. Oxford University Press, 34(10), pp. 2100–2103.
19. Omar, S. H. and Al-Wabel, N. A. (2010) 'Organosulfur compounds and possible mechanism of garlic in cancer', *Saudi Pharmaceutical Journal*. Elsevier, 18(1),

- pp. 51–58.
20. Oosthuizen, C. B., Reid, A.-M. and Lall, N.(2018) 'Garlic (*Allium sativum*) and Its Associated Molecules, as Medicine', in *Medicinal Plants for Holistic Health and WellBeing*. Elsevier, pp. 277–295.
 21. Augusti, K. T. (1996) 'Therapeutic values of onion (*Allium cepa* L.) and garlic (*Allium sativum* L.)', *Indian journal of experimental biology*, 34(7), pp. 634–640.
 22. Krishnaveni, M. et al. (2010) 'Antidiabetic and antihyperlipidemic properties of *Phyllanthus emblica* Linn. (Euphorbiaceae) on streptozotocin induced diabetic rats', *Pak J Nutr*, 9(1), pp. 43–51.
 23. Akinyemiju, T. et al. (2017) 'the burden of primary liver cancer and underlying etiologies from 1990 to 2015 at the global, regional, and national level: results from the Global Burden of Disease Study 2015', *JAMA oncology*. American Medical Association, 3(12), pp. 1683–1691.
 24. Hosseini, A. and Hosseinzadeh, H. (2015) 'A review on the effects of *Allium sativum* (Garlic) in metabolic syndrome', *Journal of Endocrinological Investigation*. Springer, 38(11), pp. 1147–1157.
 25. Abushofa, F. A., Bell, D. R. and Dyer, P. S. (2015) 'Mechanism of action of liver growth induced by non-genotoxic carcinogens (peroxisome proliferators)'.
 26. Reitman S, and Frankel A. (1957). Colorimetric method for determination of serum glutamate oxaloacetate and glutamic pyruvate transaminase. *Amer J Clin Pathol.*, 28: 56-58.
 27. Kind PRN, King EJ, Varley H, Gowenlock AH, Bell M. (1980). Method of practical clinical biochemistry. Heinman, London, , pp. 899-900.
 28. Dangerfield WG and Finlayson R. (1953). Estimation of bilirubin in serum. *J Clin Pathol.*, 6(3):173-177
 29. Rasgele, P. and Kaymak F. (2013). Effects of food preservative natamycin on liver enzymes and total protein in *Mus Musculus*. *Bulgarian J Agric Sci.*, 19(2): 298-302.
 30. Drotman, R. B., & Lawhorn, G. T. (1978). Serum enzymes as indicators of chemically induced liver damage. *Drug and chemical toxicology*, 1(2), 163-171.
 31. AL-Shinnawy, M. S. (2009). "Physiological effect of a food additive on some haematological and biochemical parameters of male albino rats." *Egypt. Acad. J. Biol. Sci* 2(1): 143-151.
 32. Muenchen, H. J. and Pienta, K. J. (1999) 'The role of the nuclear matrix in cancer chemotherapy', *Critical ReviewsTM in Eukaryotic Gene Expression*. Begel House Inc., 9(3–4).
 33. Ledda-Columbano, G. M. et al. (1998) 'In vivo hepatocyte proliferation is inducible through a TNF and IL-6-independent pathway', *Oncogene*. Nature Publishing Group, 17(8), p.1039.
 34. Bursch, W. et al. (1984) 'Controlled death (apoptosis) of normal and putative preneoplastic cells in rat liver following withdrawal of tumor promoters', *Carcinogenesis*. Oxford University Press, 5(4), pp. 453–458.
 35. Harris, J. C. et al. (2001) 'Antimicrobial properties of *Allium sativum* (garlic)', *Applied microbiology and biotechnology*. Springer, 57(3), pp. 282–286.
 36. Makowska, J. M. et al. (1990) 'Characterization of the hepatic responses to the short-term administration of ciprofibrate in several rat strains: Co-induction of microsomal cytochrome P-450 IVA1 and peroxisome proliferation', *Biochemical pharmacology*. Elsevier, 40(5), pp. 1083–1093.
 37. Ohaeri, O. C. (2001) 'Effect of garlic oil on the levels of various enzymes in the serum and tissue of streptozotocin diabetic rats', *Bioscience reports*. Springer, 21(1), pp. 19–24.
 38. Albano, E. (2006) 'Alcohol, oxidative stress and free radical damage', *Proceedings of the nutrition society*. Cambridge University Press, 65(3), pp. 278–290.
 39. Nasr, T., Bondock, S. and Youns, M. (2014) 'Anticancer activity of new coumarin substituted hydrazide-hydrazone derivatives', *European journal of medicinal chemistry*. Elsevier, 76, pp. 539–548.

40. Asdaq, S. M. and Inamdar, M. N. (2010) 'Potential of garlic and its active constituent, S-allyl cysteine, as antihypertensive and cardioprotective in Hepatoprotective Effect of Garlic Aqueous Extract against Hepatotoxicity Induced By Ciprofibrate in Male Albino Rats presence of captopril', *Phytomedicine*. Elsevier, 17(13), pp. 1016–1026.
41. Kroemer, G. et al. (2009) 'Classification of cell death: recommendations of the Nomenclature Committee on Cell Death 2009', *Cell death and differentiation*. Nature Publishing Group, 16(1), p. 3.
42. Al Kholafi, A. et al. (2008) 'Species-specific kinetics and zonation of hepatic DNA synthesis induced by ligands of PPAR α ', *Toxicological sciences*. Oxford University Press, 104(1), pp. 74–85.
43. Suzanne, M. et al. (1984) 'Midzonal necrosis as a pattern of hepatocellular injury after shock', *Gastroenterology*. Elsevier, 86(4), pp. 627–631.
44. Allen, J. W., Khetani, S. R. and Bhatia, S. N. (2004) 'In vitro zonation and toxicity in a hepatocyte bioreactor', *Toxicological sciences*. Oxford University Press, 84(1), pp. 110–119.
45. Pratibha, R. et al. (2006) 'Enzymatic studies of cisplatin induced oxidative stress in hepatic tissue of rats', *European journal of pharmacology*. Elsevier, 532(3), pp. 290–293.
46. Chandra-Kuntal, K., Lee, J. and Singh, S. V. (2013) 'Critical role for reactive oxygen species in apoptosis induction and cell migration inhibition by diallyl trisulfide, a cancer chemopreventive component of garlic', *Breast cancer research and treatment*. Springer, 138(1), pp. 69–79.