

Review Article

Child Malnutrition In Africa: Prevalence, Clinical Forms, Aetiology And Epidemiology.

Pascal Sengbé Koïvogui¹, Natia Joseph Kouadio², Mamadou Oury Diallo¹, Lanan Wassy Soromou¹, Abdoulaye Oury Barry¹ and Eric Ghigo^{1*}

¹ Training and Research Unit BioHafia, Department of Biology, Faculty of Science and Technology, University of Labé, BP 210, Labé, Republic of Guinea.

² Laboratory of Biotechnology Agriculture and Valorization of Biological Resources, UFR Biosciences, Felix HOUFHOÛT-BOIGNY University, 22 BP 582 Abidjan 22, Côte d'Ivoire.

Short Title: Child Malnutrition: Current Status

Abstract

Malnutrition among children is a devastating humanitarian disaster, particularly in sub-Saharan Africa. In this region, almost half of all children never reach their fifth birthday, falling silent victims to this scourge. For example, in Guinea, a third of young people suffer irreversible developmental delays, while 9% are extremely underweight. Diseases such as marasmus and kwashiorkor, known since the nineteenth century, continue to cause damage. Worldwide, 144 million children have their development threatened, and 47 million are severely undernourished. This paradox is particularly striking given that at the same time, 38 million other children are overweight or obese. The reasons for malnutrition are numerous and interrelated (food shortages, frequent infections, poor eating habits and poverty). The consequences of malnutrition are catastrophic, as it leads to increased infant mortality, cognitive problems and lasting physical sequelae. Malnutrition is identified and characterised by measuring parameters such as weight, height, clinical examinations and biological analyses. Children who suffer from malnutrition are at increased risk of falling ill (diarrhoea, pneumonia, malaria), which in turn increases the risk of mortality. Malnutrition can be treated through the use of ready-to-use therapeutic foods. Despite slow progress, the fight against child malnutrition remains a global priority. It requires a coordinated and comprehensive approach to improve the health and well-being of all children.

Keywords : child malnutrition, kwashiorkor, food supplement, Africa, Republic of Guinea.

INTRODUCTION

The fight against malnutrition represents a global health priority, explicitly integrated into the United Nations Sustainable Development Goals. However, childhood malnutrition remains a predominant public health issue in low-income countries, particularly in sub-Saharan Africa (1, 2). This region concentrates a significant portion of the global burden of undernutrition (3, 4). It was highlighted in 2024 by the World Health Organisation (WHO) that malnutrition contributes to 50% of deaths in children under five years of age in low- and middle-income countries. In the Republic of Guinea, malnutrition is a major issue. According to the Demographic and Health Survey (DHS) published in 2018, it appears that 30%

of children under five years of age suffer from chronic malnutrition, and 13% of them have a severe form of malnutrition. It should be noted that 9% of this population is affected by acute malnutrition (wasting). It has also been reported that 16% of children have mild underweight, and 5% of them suffer from a severe form of it (5). In this context, the Labé region presents particularly high prevalence rates of severe acute malnutrition: 13.9% according to the weight-for-age (W/A) index, 2.7% according to the weight-for-height (W/H) index, and 4.5% based on mid-upper arm circumference (5). The objective of the present review is to conduct an assessment of childhood malnutrition in Africa and its management.

***Corresponding Author:** Ghigo Eric, Training and Research Unit BioHafia, Department of Biology, Faculty of Science and Technology, University of Labé, BP 210, Labé, Republic of Guinea. **Email:** eric.ghigo@univ-labe.edu.gn

Received: 23-Sep-2025, Manuscript No. AITM-5145 ; **Editor Assigned:** 24-Sep-2025 ; **Reviewed:** 08-Oct-2025, QC No. AITM-5145 ; **Published:** 15-Oct-2025, DOI: 10.52338/aitm.2025.5145

Citation: Ghigo Eric. Child Malnutrition In Africa: Prevalence, Clinical Forms, Aetiology And Epidemiology. Advances in Tropical Medicine. 2025 October; 13(1). doi: 10.52338/aitm.2025.5145.

Copyright © 2025 Ghigo Eric. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

HISTORY OF MALNUTRITION

Malnutrition, as both a concept and a clinical manifestation, has distant historical antecedents. Descriptions consistent with oedematous states, diarrhoea, muscle wasting, apathy and skin alterations related to nutritional deficiency can be identified in ancient texts, including biblical writings. Medically, several forms of malnutrition exist. Marasmus, recognised as another severe form of malnutrition, constitutes a state of extreme wasting associated with a global deficiency in calories and protein. Kwashiorkor, whose first formal clinical descriptions of signs attributable to kwashiorkor were reported by Dr Fernando Hinojosa, a Mexican physician, and Léon Coindet, a French physician, published in the first issue of the "Gaceta Médica de México" under the title "Apuntes sobre una enfermedad del pueblo de la Magdalena" in 1865 (6, 7). Their observations, conducted in a Mexican village, concerned recurrent oedema in malnourished children, particularly during weaning or intercurrent illnesses. However, it was not until much later that kwashiorkor was specifically described by Cicely Delphine Williams. In 1933, while working in Africa on the Gold Coast (now Ghana), she observed weaned young children suffering symptoms such as oedema, diarrhoea, muscle wasting, apathy and skin alterations. She gave the disease its Ga name "kwashiorkor", meaning "disease of the displaced child". Kwashiorkor is a severe form of paediatric protein-energy malnutrition (7, 8). The mixed form known as "marasmic kwashiorkor" was later identified as the coexistence of the two clinical presentations (marasmus and kwashiorkor), in which the child exhibits both the oedema of kwashiorkor and the extreme wasting of marasmus. In 1926, Dr Léon Normet, a French physician, took particular interest in the effects of malnutrition and its clinical manifestations, especially in colonial territories such as Indochina. He described "Annamese bloat", a form of protein-energy malnutrition affecting young children. This condition was characterised by oedema and a distended abdomen, observed mainly in recently weaned children whose diet, centred almost exclusively on rice, was very low in protein. His work contributed to the understanding of infantile malnutrition in the early decades of the twentieth century (8-10). Today, understanding of malnutrition encompasses both macro- and micronutrient deficiencies and excesses in the lives of children under five years of age.

International awareness of the scale and consequences of malnutrition led to the creation of the Food and Agriculture Organization of the United Nations (FAO) on 16 October 1945 in Québec and the adoption of the Constitution of the World Health Organization (WHO) on 22 July 1946 in New York, which came into force on 7 April 1948. These institutions aim to promote fair and equitable health, improve nutritional status, and enhance agricultural accessibility and productivity for all

in a better world. Malnutrition also found legal expression in the Universal Declaration of Human Rights (1948). Article 25 of this text stipulates that "everyone has the right to a standard of living adequate for the health and well-being of himself and of his family, including food".

CURRENT GLOBAL MALNUTRITION SITUATION

Worldwide, over 144 million children under five years of age are estimated to suffer from stunting (low height for age) and 47 million from wasting (low weight for height). Concurrently, 38.3 million children are overweight or obese. Under-nutrition, whether occurring in isolation or alongside obesity, defines the "double burden of malnutrition" and is implicated in 2.7 million childhood deaths annually, representing 45% of total mortality in this age group. Each year, 6.3 million children under five die globally, nearly half of these deaths attributable to under-nutrition. Sub-Saharan Africa is among the most severely affected regions (3, 4). It is estimated that 160 million children suffer stunting on the African continent, with particularly high prevalence in West, Central and East Africa (11). Moreover, according to the 2024–2025 Global Report on Food Crises, over 36 million children across 32 countries suffer acute malnutrition, including kwashiorkor. Severe acute malnutrition without oedema (severe wasting) affects millions of African children, with around 4.5 million severely wasted children receiving treatment in 2018 according to the WHO (12-15).

In 2018, DHS data indicate that 30% of Guinean children under five years of age suffer stunting or chronic malnutrition (low height for age), 13% with severe stunting, 9% with moderate wasting (low weight for height) and 4% with severe wasting, while 16% have moderate underweight (low weight for age) and 5% severe underweight (5). Thanks to efforts and policies implemented, there has been a 9% reduction in the overall malnutrition rate, including a 4% reduction in severe acute malnutrition, according to the 2022 SMART survey (16). The Labé region in the Republic of Guinea faces a concerning situation of infant malnutrition, particularly among children aged 6 to 24 months. Despite abundant resources, malnutrition remains significant: 7.4% of children suffer acute malnutrition, characterised by rapid weight loss and general health deterioration, with 1.5% affected by severe malnutrition. Chronic malnutrition, manifesting as stunting, affects 43.3% of children, leading to physical and mental developmental issues (17, 18).

ASSESSING MALNUTRITION

Since 1995, several Malnutrition Screening Tools (MST) have been developed and implemented for early detection and classification of malnutrition forms. These techniques, often

complex and graduated, incorporate parameters beyond growth indicators alone (19, 20). Among the MSTs developed are: the Paediatric Nutritional Risk Score (PNRS), introduced in 2000 (21); the Paediatric Yorkhill Malnutrition Screening (PYMS) in 2011, comprising four stages: body mass index (BMI) below -2 standard deviations, weight-loss history, change in nutritional intake and health status over the previous week (22); and the Screening Tool for the Assessment of Malnutrition in Paediatrics (STAMP) in 2012, which assesses health status, nutritional intake and anthropometric data (23). From 2015 onwards, new tools were introduced for the general paediatric population: the Paediatric Digital Scaled Malnutrition Risk Screening Tool (PeDiSMART) and the Paediatric Nutrition Screening Tool (PNST), both evaluating health status, nutritional intake and anthropometric data (23); and the Paediatric Malnutrition Screening Tool (PMST), derived from STAMP, evaluating three main criteria: presence of a nutritionally relevant diagnosis (including overweight and obesity), changes or reduction in nutritional intake, and classification of weight and height according to percentiles or standard deviations (19, 20, 24, 25). These tools enable healthcare professionals to detect malnutrition earlier and to classify children according to the progression of nutritional complications. However, inadequate use of these tools, due to lack of proficiency, information and training of health personnel, constitutes an additional challenge in developing countries. It should be noted that the WHO relies mainly on anthropometric measures (weight, height, BMI) and the international growth reference of the National Center for Health Statistics (NCHS)/WHO (26, 27).

CAUSES OF CHILD MALNUTRITION

The aetiology of malnutrition is multifactorial; a single cause can induce multiple effects, and vice versa, an effect can influence an individual's nutritional status, leading to immediate and long-term consequences (28). The three immediate determinants of child growth—diet, health status (disease) and care practices—are closely interdependent and influenced by sociocultural, economic or climate change factors (28-30). Three types of causes of malnutrition exist: direct or immediate causes, underlying causes and fundamental or structural causes. Direct or immediate causes result from the individual immune system's response to inadequate dietary intake and/or infectious diseases (31, 32). Analysis of malnutrition determinants reveals an interlocking of factors operating at different levels. At the household and community level, three main dimensions are traditionally distinguished. First, food insecurity, characterised by limited, unstable or inadequate access to the necessary food resources. Second, child feeding and care practices, often influenced by social norms, time constraints or a lack of nutritional knowledge.

Third, public health deficiencies, including restricted access to healthcare services, sanitation infrastructure, safe water and favourable environmental conditions (33, 34). At a more global level, fundamental or structural causes relate to the socio-economic and political dynamics that frame society as a whole. These include unequal distribution of resources and financial capital, which generates disparities in access and increases the vulnerability of the most fragile groups (31, 32). Thus, malnutrition cannot be apprehended solely as the result of individual behaviours or local conditions, but must be situated within a complex system where micro- and macro-structural factors interact.

CONSEQUENCES OF CHILD MALNUTRITION

Malnutrition, defined as an imbalance resulting from insufficient or excessive intake of one or more essential nutrients, causes significant detrimental consequences for both the individual and the community. This situation contributes to stunting, affecting about one-third of children under five in these contexts (2). Children under five are particularly vulnerable to the effects of malnutrition due to their specific nutritional needs, essential for their physical, intellectual and mental development (35). Malnutrition is associated with increased infant mortality, a higher risk of chronic diseases and cognitive impairments (36). Stunting has long-term effects on affected individuals, notably in terms of cognition, educational attainment, economic productivity in adulthood and reproductive health (37, 38). The main causes of stunting include maternal nutrition and foetal growth (intrauterine growth restriction) (29).

CLINICAL FORMS OF CHILD MALNUTRITION

A child's nutritional status is modulated by numerous direct and indirect factors, including care practices, food accessibility and availability, living environment, and socio-economic and cultural status (33, 39-43). Child malnutrition manifests in various forms, notably under-nutrition, micronutrient deficiencies (hidden hunger), overweight and obesity. Clinically, these states may present as wasting, stunting, pallor, the appearance of oedema and increased susceptibility to infection. Each manifestation severely impacts a child's health and development. Frequent infections—such as measles, diarrhoea, respiratory infections and malaria—can precipitate or exacerbate acute malnutrition (44, 45). The current phenotypic classification of malnutrition, based on anthropometric measurements or oedema presence, does not systematically reflect the underlying aetiology or specific nutritional deficits (1). Infant mortality is greatly increased when clinical indications of stunting and wasting occur (46, 47). Therefore, a multidisciplinary strategy that incorporates many

indicator types to present an accurate picture of nutritional status is necessary for the assessment of child malnutrition. To assess the degree and type of malnutrition, medical practitioners employ a variety of metrics rather than relying solely on one, which is essential for efficient treatment (48). Assessment of child malnutrition generally relies on three indicator types: anthropometric, biochemical and clinical. These tools are essential to determine the degree of malnutrition severity (48). Three main types are identified: severe acute malnutrition (SAM), chronic malnutrition manifesting as stunting, and underweight. Severe acute malnutrition is characterised by rapid and significant weight loss, leading to a weight-for-height index well below age-appropriate standards. Two principal clinical forms are distinguished: marasmus, defined by extreme muscle atrophy, a cachectic appearance, wrinkled skin and lethargy; the child often exhibits irritability and anorexia, and kwashiorkor, marked by bilateral oedema, hepatomegaly, cutaneous lesions (pellagroid dermatosis), depigmentation and hair fragility, as well as pronounced apathy (8, 32, 49, 50). Chronic malnutrition (stunting) manifests as a low height-for-age ratio, a consequence of chronic or recurrent under-nutrition, often multifactorial (26, 51). Finally, underweight refers to children with low weight for age, reflecting the combined effects of acute and chronic malnutrition (32, 41, 49, 52).

EPIDEMIOLOGY OF CHILD MALNUTRITION

Nutritional epidemiology aims to elucidate the distribution and determinants of nutritional pathologies within human populations, based on experimental and observational study findings (53, 54). Epidemiologists assess the impact of dietary intake on disease incidence by collecting data from large cohorts and comparing morbidity incidence between population subgroups. Statistical approaches are employed to quantify the impact of a given exposure on morbidity risk within a population (54). Randomised controlled trials are considered the gold standard methodology for establishing causal links (53). Nevertheless, other techniques are also employed for diagnosis, such as food frequency questionnaires (e.g. 24-hour recalls) and biomarker measurements, including serum proteins (albumin, transthyretin, transferrin, C-reactive protein), electrolytes and minerals (sodium, potassium, calcium, phosphorus, magnesium), micronutrients (iron as ferritin and total iron-binding capacity, zinc, selenium, vitamin A, vitamins B12, D and C), anaemia markers (haemoglobin, haematocrit, complete blood count), inflammatory parameters (C-reactive protein, erythrocyte sedimentation rate), metabolic markers (glucose, plasma lipids such as cholesterol and triglycerides), and context-specific markers (thyroid hormones, cortisol, leptin, insulin to assess energy metabolism). Each of these tools allows interpretation of

epidemiological malnutrition data, which must imperatively consider the physiological and metabolic particularities of malnourished children and their increased risk of repeated infections and mortality (1, 22, 55-60).

Several pathologies play a major role in the onset or aggravation of child malnutrition. A self-perpetuating negative feedback loop results from the complicated, reciprocal interplay between infection and starvation. The main causes of this nutritional deterioration are frequent or persistent infections, including measles, respiratory ailments, diarrhea, and malaria. Numerous interconnected physiological pathways are to blame for this. First, the body uses a lot more energy when the immune system reacts to a disease, which raises the need for nutrition. At the same time, anorexia can be brought on by infection symptoms like fever and systemic inflammation, which results in a decreased intake of vital foods. This interaction frequently centers on the gastrointestinal tract; infections have the ability to directly harm the intestinal mucosa, which hinders the absorption of essential nutrients. In disorders such cystic fibrosis, inflammatory bowel disease (IBD), and enteropathies, where tissue damage or enzymatic malfunction impairs the absorption of vitamins, minerals, and macronutrients, this decreased absorption is also a basic feature. Consequently, this state of malnutrition compromises the host's immune system, making them more susceptible to subsequent infections and prolonging existing ones. This escalating relationship, where disease fuels malnutrition and malnutrition, in turn, exacerbates disease, presents a significant public health challenge. Metabolic and endocrine pathologies such as congenital hypothyroidism, diabetes or certain mitochondrial diseases also contribute. Some congenital or genetic anomalies, for example malabsorption syndromes or gastrointestinal tract malformations, also predispose to malnutrition. In addition, eating disorders, neglect or abuse contribute to this state. Finally, socio-economic and environmental factors such as poverty, food insecurity and poor hygiene conditions facilitate repeated infections and worsen malnutrition (34, 49, 61-63).

Malnourished children are particularly vulnerable to several severe infections, notably diarrhoea, acute respiratory infections such as pneumonia, malaria and measles. They may also suffer from severe opportunistic infections such as sepsis, septic shock, meningitis, as well as urinary infections and sometimes tuberculosis. HIV infection further exacerbates malnutrition by causing profound immunodepression. Malnutrition weakens the immune system, making infections more frequent, prolonged and severe. In turn, these infections increase energy demands, diminish appetite and disrupt nutrient absorption, creating a vicious cycle between malnutrition and infection that heightens morbidity and mortality in children. This phenomenon explains why over 70% of deaths in children under five are associated with the

malnutrition–infection combination. Specialised and tailored care is essential to reduce these risks and improve survival of malnourished children (1, 2, 64-66).

MANAGEMENT OF CHILD MALNUTRITION

Management is based on a systemic approach encompassing screening, triage and treatment of children, stratified by nutritional status severity. It utilises specialised nutritional preparations or ready-to-use therapeutic foods (RUTF), which are nutrient-dense pastes designed to treat severe acute malnutrition in children, composed mainly of peanut paste, powdered milk, vegetable oils, vitamins and minerals. Their ready-to-eat form facilitates outpatient use, often avoiding hospitalisation. The most commonly used preparations for managing severely malnourished children are F100 milk and RUTF supplemented with F75 (RUTF + F75) (34, 67-69). In complicated SAM cases, management within hospital settings is recommended (69-71). Children with uncomplicated severe acute malnutrition have increased energy needs to allow catch-up growth and weight recovery (67, 72, 73). Lyophilised supplements are also widely used, such as lipid-based nutrient supplements (LNS), fortified blended flours (FBF) and fibre-based supplements to improve anthropometric recovery; these are used for moderate acute malnutrition management (74, 75). In any case, use of RUTF or supplements requires dietary counselling and appropriate medical treatment. They may be administered in hospital or on an outpatient basis, depending on malnutrition severity and clinical complications (76-79). Despite the effectiveness of RUTF in treating moderate acute malnutrition, these foods are not always culturally appropriate, as their standardised ingredients may not align with local tastes, dietary habits and resources. Their uncooked form and texture may also seem unusual in cultures where foods are traditionally heated or prepared. Furthermore, standard formulations do not always account for local diets, which may limit social acceptance, children's appetite, and thus treatment efficacy. It is therefore recommended to adapt RUTF formulations by incorporating local ingredients and familiar flavours to improve acceptability, consumption and therapeutic success in different cultural contexts. This approach aims to reconcile nutritional efficacy with respect for popular food practices to optimise the fight against child malnutrition (80-82). Thus, complementary and therapeutic foods can be replaced by other legumes or cereals according to local availability, cost and acceptability, while maintaining caloric requirements. The efficacy of this protocol has been demonstrated, although limitations have been observed in some developing countries (73, 83-85).

CONCLUSION

Establishing optimal health in adulthood is intrinsically linked to adequate childhood nutrition. Malnutrition is among the most significant risk factors for child health. The fight against malnutrition is therefore a priority for many institutions, such as the United Nations Children's Fund (UNICEF), the World Health Organization (WHO) and the Food and Agriculture Organization of the United Nations (FAO). Significant advances have been made, but current progress remains far too slow and profoundly unequal, as indicated in the 2020 Global Nutrition Report (86). Numerous obstacles—such as failing agricultural systems, social inequalities, limited access to safe water and basic healthcare, climate change and armed conflicts—continue to impede progress. Although various dietary formulations for malnutrition management currently exist, their accessibility to families remains limited. It is therefore crucial to develop and evaluate the acceptability of local products for managing malnourished children. The challenges to eradicating hunger in all its forms and achieving universal food security remain immense.

Funding

Pascal Sengbé Koïvogui, Mamadou Oury Diallo, Lanan Wassy Soromou, Abdoulaye Oury Barry and Eric Ghigo are employed by the Ministry of Higher Education, Scientific Research and Innovation (MESRSI).

Declaration Of Conflict Of Interest

The authors declare no conflicts of interest.

Author Contributions

Pascal Sengbé Koïvogui, Natia Joseph Kouadio, Mamadou Oury Diallo, Lanan Wassy Soromou, Abdoulaye Oury Barry and Eric Ghigo conducted the bibliographic research, analysed the bibliographic data, and wrote and edited the manuscript.

REFERENCES

1. Bhutta ZA, Berkley JA, Bandsma RHJ, Kerac M, Trehan I, Briend A. Severe childhood malnutrition. *Nature Reviews Disease Primers*. 2017;3(1):1-18.
2. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet*. 2013;382(9890):427-51.
3. Maphosa TP, Mulqueeny DM, Osei E, Kuupiel D, Mashamba-Thompson TP. Mapping evidence on malnutrition screening tools for children under 5 years in sub-Saharan Africa: a scoping review protocol. *Systematic Reviews*. 2020;9(1):52.

4. Agho KE, Akombi BJ, Ferdous AJ, Mbugua I, Kamara JK. Childhood undernutrition in three disadvantaged East African Districts: a multinomial analysis. *BMC Pediatrics*. 2019;19(1):118.
5. Institut National de la Statistique Ministère du Plan et du Développement Economique Conakry G. Enquête Démographique et de Santé en Guinée 2018 : Indicateurs Clés. RÉPUBLIQUE DE GUINÉE; 2018. p. 1-61.
6. Noel A, Morel FB, Marion-Letellier R. La malnutrition aiguë sévère : de la physiopathologie à la prise en charge. *Nutrition Clinique et Métabolisme*. 2024;38(1):2-10.
7. Benjamin O, Lappin SL. Kwashiorkor(Archived). StatPearls. Treasure Island (FL) ineligible companies. Disclosure: Sarah Lappin declares no relevant financial relationships with ineligible companies.2025.
8. Heikens GT, Manary M. 75 years of Kwashiorkor in Africa. *Malawi Med J*. 2009;21(3):96-8.
9. Nott J. "No one may starve in the British Empire": Kwashiorkor, Protein and the Politics of Nutrition Between Britain and Africa. *Soc Hist Med*. 2021;34(2):553-76.
10. Pham TP, Alou MT, Golden MH, Million M, Raoult D. Difference between kwashiorkor and marasmus: Comparative meta-analysis of pathogenic characteristics and implications for treatment. *Microb Pathog*. 2021;150:104702.
11. Simwanza NR, Kalungwe M, Karonga T, Mtambo CMM, Ekpenyong MS, Nyashanu M. Exploring the risk factors of child malnutrition in Sub-Sahara Africa: A scoping review. *Nutrition and Health*. 2022.
12. Wambua J, Ali A, Ukwizabigira JB, Kuodi P. Prevalence and risk factors of under-five mortality due to severe acute malnutrition in Africa: a systematic review and meta-analysis. *Syst Rev*. 2025;14(1):29.
13. Munthali T, Jacobs C, Sitali L, Dambe R, Michelo C. Mortality and morbidity patterns in under-five children with severe acute malnutrition (SAM) in Zambia: a five-year retrospective review of hospital-based records (2009–2013). *Archives of Public Health*. 2015;73(1):23.
14. Williams PCM, Berkley JA. Guidelines for the treatment of severe acute malnutrition: a systematic review of the evidence for antimicrobial therapy. *Paediatr Int Child Health*. 2018;38(sup1):S32-S49.
15. Carboo JA, Lombard M, Conradie C, Dolman RC, Ricci C. Evaluation of the treatment guidelines, practices and outcomes of complicated severe acute malnutrition in children aged 0-59 months in sub-Saharan Africa: a study protocol for the SAMAC study. *Pan Afr Med J*. 2020;36:241.
16. Guinée U. enquête nutritionnelle SMART 2022 Guinée. 2023.
17. Guinée U. Pour l'année 2022, près de 200000 enfants seraient exposés à la malnutrition aiguë sévère en Guinée 2023 [updated 2023].
18. Camara Mm, Tall I, Daffe M. Concertation Regionale Sur La Situation Alimentaire Et Nutritionnelle Au Sahel Et En Afrique De L'ouest Dakar (Senegal), 22-24 Mars 2017 2017.
19. McCarthy A, Delvin E, Marcil V, Belanger V, Marchand V, Boctor D, et al. Prevalence of Malnutrition in Pediatric Hospitals in Developed and In-Transition Countries: The Impact of Hospital Practices. *Nutrients*. 2019;11(2):236.
20. Huysentruyt K, Vandenplas Y, De Schepper J. Screening and assessment tools for pediatric malnutrition. *Current Opinion in Clinical Nutrition & Metabolic Care*. 2016;19(5):336.
21. Joosten K, van der Velde K, Joosten P, Rutten H, Hulst J, Dulfer K. Association between nutritional status and subjective health status in chronically ill children attending special schools. *Quality of Life Research*. 2016;25(4):969-77.
22. McNamara AE, Brennan L. Potential of food intake biomarkers in nutrition research. *Proceedings of the Nutrition Society*. 2020;79(4):487-97.
23. Ong SH, Chee WSS, Lapchmanan LM, Ong SN, Lua ZC, Yeo JX-N. Validation of the Subjective Global Nutrition Assessment (SGNA) and Screening Tool for the Assessment of Malnutrition in Paediatrics (STAMP) to Identify Malnutrition in Hospitalized Malaysian Children. *Journal of Tropical Pediatrics*. 2019;65(1):39-45.
24. Sayed S, El-Shabrawi MHF, Abdelmonaem E, El Koofy N, Tarek S. Value of Nutritional Screening Tools Versus Anthropometric Measurements in Evaluating Nutritional Status of Children in a Low/Middle-Income Country. *Pediatric Gastroenterology, Hepatology & Nutrition*. 2023;26(4):213-23.

25. Nel E. Severe acute malnutrition. *Current Opinion in Clinical Nutrition & Metabolic Care*. 2018;21(3):195.
26. Onis Md, Onyango AW, Borghi E, Garza C, Yang H, Group WMGRS. Comparison of the World Health Organization (WHO) Child Growth Standards and the National Center for Health Statistics/WHO international growth reference: implications for child health programmes. *Public Health Nutrition*. 2006;9(7):942-7.
27. Ackatia-Armah RS, McDonald CM, Doumbia S, Erhardt JG, Hamer DH, Brown KH. Les enfants maliens souffrant de malnutrition aiguë modérée qui sont traités avec des compléments alimentaires à base de lipides ont des gains de poids et des taux de récupération plus élevés que ceux traités avec des produits à base de céréales et de légumineuses produits localement : un essai communautaire randomisé en grappes. *The American Journal of Clinical Nutrition*. 2015;101(3):632-45.
28. Fan Y, Yao Q, Liu Y, Jia T, Zhang J, Jiang E. Underlying Causes and Co-existence of Malnutrition and Infections: An Exceedingly Common Death Risk in Cancer. *Front Nutr*. 2022;9:814095.
29. Frongillo EA, Jr. Symposium: Causes and Etiology of Stunting. Introduction. *J Nutr*. 1999;129(2S Suppl):529S-30S.
30. Romanello M, Napoli Cd, Green C, Kennard H, Lampard P, Scamman D, et al. The 2023 report of the Lancet Countdown on health and climate change: the imperative for a health-centred response in a world facing irreversible harms. *Lancet (London, England)*. 2023;402(10419):2346-94.
31. Grover Z, MBBS. Malnutrition protéino-énergétique - Cliniques pédiatriques. 2009.
32. Koletzko B, Bhutta ZA, Cai W, Dhansay MA, Duggan CP, Makrides M, et al. *Pediatric Nutrition in Practice*: Karger Medical and Scientific Publishers; 2022 2022-03.
33. Ahmed. Malnutrition - Ahmed - 2012 - Journal de gastroentérologie et de nutrition pédiatriques - Bibliothèque en ligne Wiley 2012 [updated 2012].
34. Dipasquale V, Cucinotta U, Romano C. Acute Malnutrition in Children: Pathophysiology, Clinical Effects and Treatment. *Nutrients*. 2020;12(8):2413.
35. Govender I, Rangiah S, Kaswa R, Nzaumvila D. Malnutrition in children under the age of 5 years in a primary health care setting. *S Afr Fam Pract* (2004). 2021;63(1):e1-e6.
36. Huffman SL, Schofield D. Consequences of malnutrition in early life and strategies to improve maternal and child diets through targeted fortified products. *Maternal & Child Nutrition*. 2011;7(s3):1-4.
37. Dewey KG, Begum K. Long-term consequences of stunting in early life. *Maternal & Child Nutrition*. 2011;7 Suppl 3(Suppl 3):5-18.
38. Bosch AM, Baqui AH, van Ginneken JK. Early-life determinants of stunted adolescent girls and boys in Matlab, Bangladesh. *Journal of Health, Population, and Nutrition*. 2008;26(2):189-99.
39. Bhutta. Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost? *The Lancet*. 2013;382(9890):452-77.
40. Corsi DJ, Mejía-Guevara I, Subramanian SV. Risk factors for chronic undernutrition among children in India: Estimating relative importance, population attributable risk and fractions. *Social Science & Medicine*. 2016;157:165-85.
41. Evans WJ, Morley JE, Argilés J, Bales C, Baracos V, Guttridge D, et al. Cachexia: A new definition. *Clinical Nutrition*. 2008;27(6):793-9.
42. Ruel MT, Alderman H. Nutrition-sensitive interventions and programmes: how can they help to accelerate progress in improving maternal and child nutrition? *The Lancet*. 2013;382(9891):536-51.
43. Siddiqi M, Zubair A, Kamal A, Ijaz M, Aljeddani SMA. Risk factors of child malnutrition under 5 years: Evidence from Pakistan using the Composite Index of Anthropometric Failure. *Nutrition*. 2024;127:112523.
44. Michaelsen KF, Hoppe C, Roos N, Kaestel P, Stougaard M, Lauritzen L, et al. Choice of foods and ingredients for moderately malnourished children 6 months to 5 years of age. *Food Nutr Bull*. 2009;30(3 Suppl):S343-404.
45. Rice AL, Sacco L, Hyder A, Black RE. Malnutrition as an underlying cause of childhood deaths associated with infectious diseases in developing countries. *Bull World Health Organ*. 2000;78(10):1207-21.

46. Thurstans S, Sessions N, Dolan C, Sadler K, Cichon B, Isanaka S, et al. The relationship between wasting and stunting in young children: A systematic review. *Maternal & Child Nutrition*. 2022;18(1):e13246.
47. Ando Y, Sakamoto A, Marusawa H. Unexpected anti-tumor effect of immune checkpoint inhibitors followed by transcatheter arterial chemoembolization for hepatocellular carcinoma. *Clin J Gastroenterol*. 2023;16(6):891-4.
48. Dewey KG, Cohen RJ, Rollins NC. Feeding of Nonbreastfed Children from 6 to 24 Months of Age in Developing Countries. *Food and Nutrition Bulletin*. 2004;25(4):377-402.
49. Mehta NM, Corkins MR, Lyman B, Malone A, Goday PS, Carney LN, et al. Defining Pediatric Malnutrition. *Journal of Parenteral and Enteral Nutrition*. 2013;37(4):460-81.
50. Yessoufou AG, Behanzin J, Ahokpe M, Djinti SA, Bossou R, Sezan A. Prévalence de l'anémie chez les enfants malnutris de 6 à 59 mois hospitalisés au CHD/Zou-Collines dans le plateau d'Abomey (Centre du Bénin). *International Journal of Biological and Chemical Sciences*. 2015;9(1):82-90.
51. De Sanctis V, Soliman A, Alaaraj N, Ahmed S, Alyafei F, Hamed N. Early and Long-term Consequences of Nutritional Stunting: From Childhood to Adulthood. *Acta Biomed*. 2021;92(1):e2021168.
52. Monga M, Sikorski C, de Silva H, McGrath M, Kerac M. Identifying underweight in infants and children using growth charts, lookup tables and a novel "MAMI" slide chart: A cross-over diagnostic and acceptability study. *PLOS Glob Public Health*. 2023;3(8):e0002303.
53. Sullivan VK, Rebholz CM. Nutritional Epidemiology and Dietary Assessment for Patients With Kidney Disease: A Primer. *American Journal of Kidney Diseases*. 2023;81(6):717-27.
54. Satija A, Yu E, Willett WC, Hu FB. Understanding Nutritional Epidemiology and Its Role in Policy. *Advances in Nutrition*. 2015;6(1):5-18.
55. Dhurandhar NV, Schoeller D, Brown AW, Heymsfield SB, Thomas D, Sørensen TIA, et al. Energy balance measurement: when something is not better than nothing. *International Journal of Obesity*. 2015;39(7):1109-13.
56. Golden MH. Evolution of nutritional management of acute malnutrition. *Indian Pediatrics*. 2010;47(8):667-78.
57. Kipnis V, Midthune D, Freedman L, Bingham S, Day NE, Riboli E, et al. Bias in dietary-report instruments and its implications for nutritional epidemiology. *Public Health Nutrition*. 2002;5(6a):915-23.
58. Rafiq T, Azab SM, Teo KK, Thabane L, Anand SS, Morrison KM, et al. Nutritional Metabolomics and the Classification of Dietary Biomarker Candidates: A Critical Review. *Advances in Nutrition*. 2021;12(6):2333-57.
59. Tinker LF, Sarto GE, Howard BV, Huang Y, Neuhauser ML, Mossavar-Rahmani Y, et al. Associations entre l'apport énergétique et protéique alimentaire calibré par biomarqueurs et le risque de diabète chez les femmes ménopausées, selon la Women's Health Initiative 1 2 3 4. *The American Journal of Clinical Nutrition*. 2011;94(6):1600-6.
60. Zhang Z, Pereira SL, Luo M, Matheson EM. Evaluation of Blood Biomarkers Associated with Risk of Malnutrition in Older Adults: A Systematic Review and Meta-Analysis. *Nutrients*. 2017;9(8).
61. Hendricks KM, Duggan C, Gallagher L, Carlin AC, Richardson DS, Collier SB, et al. Malnutrition in Hospitalized Pediatric Patients: Current Prevalence. *Archives of Pediatrics & Adolescent Medicine*. 1995;149(10):1118-22.
62. Isabelle MN, Annick NN, Drame H, Jeannette EN, Jocelyn TN, Selangai KH, et al. Aspects Epidémiologiques, Cliniques, Biologiques et Evolutifs de l'Anémie Sévère chez l'Enfant à Yaoundé. *HEALTH SCIENCES AND DISEASE*. 2024;25(4).
63. Pelletier DL, Frongillo EA, Habicht JP. Epidemiologic evidence for a potentiating effect of malnutrition on child mortality. *American Journal of Public Health*. 1993;83(8):1130-3.
64. Chevalier P, Delpeuch F, Maire B. [The "malnutrition-infection" complex, the most widespread public health problem in underprivileged populations.]. *Med Mal Infect*. 1996;26 Suppl 3:366-70.
65. Kambale RM, Kasengi JB, Kivukuto JM, Cubaka LM, Mungo BM, Balaluka GB. [Infectious profile and mortality of children aged 0-5 years admitted with severe acute malnutrition: a retrospective cohort study conducted in

- the Therapeutic Nutritional Center in Bukavu, Republic of Congo]. *Pan Afr Med J.* 2016;23:139.
66. Saengnipanthkul S, Phosuwattanakul J, Thepsuthammarat K, Chongviriyaphan N. Epidemiological data on nutritional disorders and outcomes in hospitalized Thai children: an analysis of data from the National Health Database 2015-2019. *Epidemiology and Health.* 2022;44:e2022047.
 67. Ashworth A, Organization WH. Guidelines for the Inpatient Treatment of Severely Malnourished Children: World Health Organization; 2003 2003. 50 p.
 68. Bailey J, Opondo C, Lelijveld N, Marron B, Onyo P, Musyoki EN, et al. A simplified, combined protocol versus standard treatment for acute malnutrition in children 6-59 months (ComPAS trial): A cluster-randomized controlled non-inferiority trial in Kenya and South Sudan. *PLoS Med.* 2020;17(7):e1003192.
 69. Versloot CJ, Voskuijl W, van Vliet SJ, van den Heuvel M, Carter JC, Phiri A, et al. Effectiveness of three commonly used transition phase diets in the inpatient management of children with severe acute malnutrition: a pilot randomized controlled trial in Malawi. *BMC Pediatrics.* 2017;17:112.
 70. Ashworth A. Efficacy and Effectiveness of Community-Based Treatment of Severe Malnutrition. *Food and Nutrition Bulletin.* 2006;27(3_suppl3):S24-S48.
 71. Akomo P, Bahwere P, Murakami H, Banda C, Maganga E, Kathumba S, et al. Soya, maize and sorghum ready-to-use therapeutic foods are more effective in correcting anaemia and iron deficiency than the standard ready-to-use therapeutic food: randomized controlled trial. *BMC Public Health.* 2019;19(1):806.
 72. James G, Stephenson K, Callespie M. Stabilité de l'acide docosahexaénoïque dans les aliments thérapeutiques prêts à l'emploi. 2023.
 73. Rachmadewi A, Soekarjo DD, Bait BR, Suryantan J, Noor R, Rah JH, et al. Ready-to-Use Therapeutic Foods (RUTFs) Based on Local Recipes Are as Efficacious and Have a Higher Acceptability than a Standard Peanut-Based RUTF: A Randomized Controlled Trial in Indonesia. *Nutrients.* 2023;15(14):3166.
 74. Lazzerini M, Rubert L, Pani P. Specially formulated foods for treating children with moderate acute malnutrition in low- and middle-income countries - Lazzerini, M - 2013 | Cochrane Library.
 75. Dewey KG, Stewart CP, Wessells KR, Prado EL, Arnold CD. Small-quantity lipid-based nutrient supplements for the prevention of child malnutrition and promotion of healthy development: overview of individual participant data meta-analysis and programmatic implications. *Am J Clin Nutr.* 2021;114(Suppl 1):3S-14S.
 76. Brown KH, Nyirandutiye DH, Jungjohann S. Management of children with acute malnutrition in resource-poor settings. *Nature Reviews Endocrinology.* 2009;5(11):597-603.
 77. Zoungrana B, Sawadogo PS, Somda NS, Tapsoba F, Tankoano A, Savadogo A. Performance et coût de la prise en charge de la malnutrition aiguë sévère avec complications à Kaya, Burkina Faso. *Pan African Medical Journal.* 2019;34(1).
 78. Eglseder D, Bauer S, Schuettengruber G. Malnutrition management and the application of medical nutrition therapy in hospitalized patients at the end of life. *J Nutr Health Aging.* 2024;28(7):100255.
 79. Malaria WA, Malnutrition Study G. Does acute malnutrition in young children increase the risk of treatment failure following artemisinin-based combination therapy? A WWARN individual patient data meta-analysis. *Lancet Glob Health.* 2024;12(4):e631-e40.
 80. Aiello I, Kounnavong S, Vinathan H, Philavong K, Luangphaxay C, Soukhavong S, et al. Short-Term Acceptability of Ready-to-Use Therapeutic Foods in Two Provinces of Lao People's Democratic Republic. *Nutrients.* 2023;15(17).
 81. Matilsky DK, Maleta K, Castleman T, Manary MJ. Supplementary feeding with fortified spreads results in higher recovery rates than with a corn/soy blend in moderately wasted children. *J Nutr.* 2009;139(4):773-8.
 82. LaGrone LN, Trehan I, Meuli GJ, Wang RJ, Thakwalakwa C, Maleta K, et al. A novel fortified blended flour, corn-soy blend "plus-plus," is not inferior to lipid-based ready-to-use supplementary foods for the treatment of moderate acute malnutrition in Malawian children. *Am J Clin Nutr.* 2012;95(1):212-9.
 83. Puoane T, Sanders D, Chopra M, Ashworth A, Strasser S, McCoy D, et al. Evaluating the clinical management

- of severely malnourished children- a study of two rural district hospitals. South African Medical Journal. 2001;91(2):137-41.
84. Samena HSC, Rasamoelison RJ, Ramananirina MZ, Rakotomahefa Narison ML, Robinson AL. Pratique de la diversification alimentaire chez les nourrissons à Antananarivo, Madagascar : relation avec leur état nutritionnel. Journal de Pédiatrie et de Puériculture. 2023;36(3):107-13.
85. Borg B, Mihrshahi S, Laillou A, Sigh S, Sok D, Peters R, et al. Development and testing of locally-produced ready-to-use therapeutic and supplementary foods (RUTFs and RUSFs) in Cambodia: lessons learned. BMC Public Health. 2019;19(1):1200.
86. Mannar MG, Micha R. Rapport sur la nutrition mondiale. Journal de Pédiatrie et de Puériculture. 2020;36(3):10-171.