

Short Review

Immunity And Endometriosis: A Short Review.

Ludek Fiala¹, Jiri Lenz², Radek Chvatal³, Vaclav Vetvicka⁴¹Institute of Sexology, Charles University Prague Psychiatric Clinic, Charles University Pilsen, Czech Republic²Faculty of Veterinary Medicine, University of Veterinary and Pharmaceutical Science Brno, Czech Republic³Department of Obstetrics and Gynaecology, Znojmo Hospital, Znojmo, Czech Republic⁴University of Louisville, Department of Pathology, Louisville, KY 40202, USA

Abstract

Endometriosis is a prevalent disorder of the female reproductive system that frequently demonstrates malignant-like characteristics, including invasive growth and recurrence. Despite extensive investigation, the precise mechanisms underlying its pathogenesis remain incompletely elucidated. Accumulating evidence indicates that dysregulation of the immune system plays a central role in disease development and progression. Alterations in both the number and function of immune cell populations, such as macrophages, T and B lymphocytes, and natural killer (NK) cells, have been consistently documented. These findings have led to the hypothesis that immunotherapy may represent a promising novel strategy for the treatment of endometriosis.

Keywords : Endometriosis; macrophages; immunity; NK cells; autoimmunity.

INTRODUCTION

Endometriosis is a chronic medical condition in which tissue resembling the lining of the uterus grows outside the uterine cavity. These ectopic growths can occur on the ovaries, fallopian tubes, the outer surface of the uterus, the pelvic peritoneum, and in rare cases, in more distant organs. Beyond its well-known association with infertility, endometriosis is also linked to stress-related effects on sexual dysfunction.¹

It is one of the leading causes of morbidity among premenopausal women. Despite decades of extensive research, the complex pathogenesis of this enigmatic disorder remains controversial.² The most widely accepted theory is that endometriosis develops as a result of retrograde menstruation and/or transplantation of shed endometrial fragments. When displaced endometrial cells implant on pelvic structures, they remain hormonally responsive and form the characteristic lesions of the disease. However, this explanation is incomplete: retrograde menstruation occurs in more than 90% of women, yet fewer than 10% develop endometriosis.³

Alternative theories have been proposed. The metastatic theory suggests that endometrial cells disseminate through lym-

phatic or vascular routes.⁴ Another hypothesis, the stem cell theory, posits that progenitor or stem-like cells may contribute to lesion formation.⁵ For a comprehensive overview of these and other proposed mechanisms, readers are referred to.⁶ Increasing attention has been directed toward the immune system, even though endometriosis is generally regarded as a steroid-sensitive condition. Evidence shows multiple facets of immune dysfunction in affected women.⁷ Endometriosis is not simply a gynecological disease but one in which immune dysregulation plays a central role. Normally, immune surveillance clears displaced endometrial cells. In endometriosis, however, this process is impaired, resulting in chronic inflammation, angiogenesis, and lesion survival.

MACROPHAGES

Macrophages are central immune effectors responsible for pathogen recognition, phagocytosis, and antigen presentation to T cells. Under physiological conditions, they also participate in regeneration of the endometrium,⁸ a process partly regulated by CD36 receptor expression and matrix metalloproteinase activation—both reduced in endometriosis.⁹

***Corresponding Author:** Vaclav Vetvicka, University of Louisville, Department of Pathology, Louisville, KY 40202, USA. **Email:** vaclav.vetvicka1@gmail.com
Received: 20-Sep-2025, Manuscript No. TJOCM-5142 ; **Editor Assigned:** 24-Sep-2025 ; **Reviewed:** 11-Oct-2025, QC No. TJOCM-5142 ; **Published:** 16-Oct-2025,
DOI: 10.52338/tjocm.2025.5142

Citation: Vaclav Vetvicka. Immunity And Endometriosis: A Short Review. The Journal of Clinical Medicine. 2025 October; 13(1). doi: 10.52338/tjocm.2025.5142.
Copyright © 2025 Vaclav Vetvicka. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The role of macrophages in endometriosis remains debated. Their numbers are consistently elevated in the peritoneal cavity, yet their functional activity appears suppressed.⁸ Reduced expression of CD3, CD36, and annexin A2 may underlie this impairment.¹⁰ Activated macrophages release proinflammatory cytokines such as TNF- α , IL-6, IL-8, and IL-1 β , promoting neutrophil recruitment, chronic inflammation, and lesion development.¹¹

Lesional macrophages often display an M2-polarized phenotype,¹² associated with tissue remodeling and cancer progression.¹³ Studies suggest a dynamic shift from M1 dominance in early disease to M2 predominance in later stages, indicating a role in disease progression.¹⁴ These findings have prompted interest in macrophage-targeted therapies as a potential treatment strategy.¹⁵

LYMPHOCYTES

T lymphocytes represent another key immune population implicated in endometriosis. CD8⁺ T cells are the predominant T-cell subset in endometrial tissue, found as single intraepithelial cells, adjacent to glands, or clustered in lymphoid aggregates.¹⁶ However, data on their activity and function remain inconsistent.¹⁷ B lymphocytes have also been studied, though results are similarly inconclusive. While many studies report increased B-cell activation and autoantibody production, others describe unchanged or reduced B-cell counts.¹⁸ The pathogenic significance of autoantibodies remains uncertain.

More recent research has examined regulatory lymphocyte subsets. Tregs and Bregs from women with endometriosis show altered numbers and enhanced secretion of IL-24, which may contribute to immunosuppression and ectopic tissue survival.¹⁹ Elevated PD-1 and PD-L1 expression on circulating T and B cells has also been documented, potentially contributing to immune tolerance of endometriotic lesions.²⁰ Interestingly, activated Tregs appear reduced within endometriotic tissue but not in peripheral blood, suggesting a localized effect on inflammation and lesion growth.²¹

NK CELLS

NK cells normally play a critical role in immune surveillance by eliminating abnormal or misplaced cells. In endometriosis, their cytotoxic function is impaired, which may allow ectopic endometrial fragments to evade clearance and persist in the peritoneal cavity.²²

Reduced NK cell activity has been consistently observed in patients with endometriosis and may serve as a potential diagnostic marker.²³ Mechanistically, the disease is associated with upregulation of NK inhibitory receptors (e.g., KIR2DL1, CD94/NKG2A, PD-1) and downregulation of activating

receptors (e.g., NKp46, NKp30, NKG2D), leading to impaired cytotoxicity.

Beyond immune evasion, NK cells may influence stromal cell biology, local immune tolerance, and implantation processes, thereby contributing to infertility and pregnancy complications. For an in-depth review, see.²⁴

CYTOKINES

Cytokine imbalance is another hallmark of endometriosis. Women with the disease exhibit elevated levels of proinflammatory and growth-related cytokines such as GM-CSF, IL-1, IL-6, IL-8, IL-10, TNF- α , IGF-1, and TGF- β .²⁵ These mediators promote adhesion, invasion, angiogenesis, and survival of ectopic lesions.

Despite clear associations, causality remains uncertain. For instance, elevated IL-1 production has been reported,²⁶ but results vary across studies. Large-scale profiling has revealed distinct cytokine expression signatures, but their mechanistic roles are poorly defined.^{27, 28} At present, cytokines may be more promising as diagnostic biomarkers than as established pathogenic drivers.²⁹ Therapeutic strategies targeting cytokines—such as TNF inhibitors, IL-6 blockers, and chemokine antagonists—are under investigation but show variable outcomes.³⁰

AUTOIMMUNITY

Endometriosis shares certain features with autoimmune disorders, including chronic inflammation, autoantibody production, and altered lymphocyte populations.^{31, 32} However, available evidence does not support its classification as a true autoimmune disease. Elevated autoantibodies typically target endometrial antigens, suggesting they are a consequence rather than a cause of disease. While epidemiological links with autoimmune and allergic diseases have been suggested, they remain unconfirmed.³³ Nonetheless, immunomodulatory strategies used in autoimmunity have been proposed as potential therapeutic approaches.³⁴

CONCLUSIONS

Endometriosis is increasingly viewed as a disorder of immune dysregulation. Substantial evidence indicates widespread alterations in the activity of immune cells—including macrophages, lymphocytes, NK cells—as well as in cytokine networks. However, it remains unclear whether these immune changes are causal factors or consequences of the disease.

While immunotherapy offers a promising avenue, its clinical application remains limited. Current immunomodulators have shown only modest effects.³⁰ Given the unsatisfactory outcomes of conventional treatments, future therapies may

combine classical hormonal approaches with novel strategies, including selective progesterone receptor modulators, aromatase inhibitors, anti-angiogenic agents, GnRH antagonists, and even gene-based interventions.³⁵ Continued investigation into the immunological mechanisms underlying endometriosis will be essential for developing more effective, targeted therapies.

REFERENCES

1. Fiala, L., Lenz, J., Bob, P. 2021. Effect of psychosocial trauma and stress on sexual dysfunction in women with endometriosis. *Medicine (Baltimore)*, 100(31):e26836. doi: 10.1097/MD.00000000000026836.
2. Baldi, A., Campioni, M., Signorile, P.G. 2008. Endometriosis: pathogenesis, diagnosis, therapy and association with cancer (review). *Oncol. Rep.*, 19:843-846.
3. D'Hooghe, T.M., Debrock, S. 2002. Endometriosis, retrograde menstruation and peritoneal inflammation in women and in baboons. *Hum. Reprod. Update*, 8:84-88.
4. Sampson, J.A. 1927. Metastatic or embolic endometriosis, due to the menstrual dissemination of endometrial tissue into the venous circulation. *Am. J. Pathol.*, 3: 93-110.
5. Yan, Z., Wang, Y., Cragun, J.M., Chambers, S.K., Zheng, W. 2013. Cell origin of endometriosis: contribution by the fallopian tube epithelium. *Am. J. Clin. Exp. Obstet. Gynecol.*, 1:37-42.
6. Lamceva, J., Uljanovs, R., Strumfa, I. 2023. The main theories on the pathogenesis of endometriosis. *Int. J. Mol. Sci.*, 24, doi:10.3390/ijms24054254
7. Kralickova, M., Vetvicka, V. 2015. Immunological aspects of endometriosis: A review. *Ann. Transl. Med.*, 3, doi: 10.3978/j.issn.2305-5839.2015.06.08
8. Abramiuk, M., Grywalska, E., Małkowska, P., Sierawska, O., Hryniewicz, R., Niedźwiedzka-Rystwej, P. 2022. The role of the immune system in the development of endometriosis. *Cells*, 11:2028, doi: 10.3390/cells11132028.
9. Wu, M.H., Shoji, Y., Wu, M.C., Chuang, P.C., Lin, C.C., Huang, M.F., Tsai, S.J. 2005. Suppression of matrix metalloproteinase-9 by prostaglandin E(2) in peritoneal macrophage is associated with severity of endometriosis. *Am. J. Pathol.*, 167:1061-1069.
10. Wu, M.H., Chuang, P.C., Lin, Y.J., Tsai, S.J. 2013. Suppression of annexin A2 by prostaglandin E₂ impairs phagocytic ability of peritoneal macrophages in women with endometriosis. *Hum. Reprod.*, 28:1045-1053.
11. Smolarz, B., Szyłło, K., Romanowicz, H. 2021. Endometriosis: Epidemiology, classification, pathogenesis, treatment and genetics (Review of literature) *Int. J. Mol. Sci.*, 22:10554, doi: 10.3390/ijms221910554.
12. Izumi, G., Koga, K., Takamura, M., Makabe, T., Satake, E., Takeuchi, A., Taguchi, A., Urata, Y., Fujii, T., Osuga, Y. 2018. Involvement of immune cells in the pathogenesis of endometriosis. *J. Obstet. Gynaecol. Res.*, 44:191-198.
13. Kralickova, M., Losan, P., Vetvicka, V. 2014. Endometriosis and cancer. *Womens Health*, 10:591-597.
14. Vallvé-Juanico, J., George, A.F., Sen, S., Thomas, R., Shin, M.-G., Kushnoor, D., Vázquez, J.J., Vo, K.C., et al., 2022. Deep immunophenotyping reveals endometriosis is marked by dysregulation of the mononuclear phagocytic system in endometrium and peripheral blood. *BMC Med.*, 20, 158 <https://doi.org/10.1186/s12916-022-02359-4>.
15. Song, L., Yang, C., Ji, G., Hu, R. 2024. The role and potential treatment of macrophages in patients with infertility and endometriosis. *J. Reprod. Immunol.*, 166: 104384.
16. Yeaman, G.R., Guyre, P.M., Fanger, M.W., Collins, J.E., White, H.D., Rathbun, W., et al. 1997. Unique CD8+ T cell-rich lymphoid aggregates in human uterine endometrium. *J. Leukoc. Biol.*, 61:427-435.
17. Kisovar, A., Becker, C.M., Granne, I., Southcombe, J.H. 2023. The role of CD8+ T cells in endometriosis: a systematic review. *Front. Immunol.*, 14:1225639, doi: 10.3389/fimmu.2023.1225639
18. Riccio, L.G.C., Barakat, E.C., Chapron, C., Batteux, F., Abrão, M.S. 2017. The role of the B lymphocytes in endometriosis: A systematic review. *J. Reprod. Immunol.*, 123: 29-34.
19. Kedzierska, A.E., Lorek, D., Slawek, A., Karmowski, M., Kalota, A., Pajak, J., Chrobak, A., Grzymajlo, K., Chelmonska-Soyta, A. 2025. IL-24 producing regulatory T and B lymphocytes in endometriosis. *Front Immunol.*, 16, doi:10.3389/fimmu.2025.1582762
20. Walankiewicz, M., Grywalska, E., Polak, G., Korona-Glowniak, I., Witt, E., Sardacka, A. 2018. The increase in circulating PD-1- and PD-L1-expressing lymphocytes in endometriosis: Correlation with clinical and laboratory parameters. *Mediators Inflamm.*, 2018: doi:10.1155/2018/7041342

21. Tanaka, Y., Mori, A.T., Ito, F., Koshiba, A., Takaoka, O, et al. 2017. Exacerbation of endometriosis due to regulatory T-cell dysfunction *J. Clin. Endocrinol. Metab.*, 102:3206-3217.
22. Thiruchelvam, U., Wingfield, M., O'Farrelly, C. 2015. Natural killer cells: Key players in endometriosis. *Am.J. Reprod. Immunol.*, 74:291–301.
23. Wang, L., Li, L., Li, Y., Huang, C., Lian, R., Wu, T., Ma, J., Zhang, Y., Cheng, Y., Diao, L., Zeng, Y. 2021. A history of endometriosis is associated with decreased peripheral NK cytotoxicity and increased infiltration of uterine CD68+ macrophages. *Front. Immunol.*, 12, 2021, doi:10.3389/fimmu.2021.711231
24. Yang, S., Wang, H., Li, D., Li, M. 2024. An estrogen–NK cells regulatory axis in endometriosis, related infertility, and miscarriage. *Int. J. Mol. Sci.*, 25: 3362, doi:10.3390/ijms25063362
25. Suryawanshi, S., Huang, X., Elishaev, E., et al. 2014. Complement pathway is frequently altered in endometriosis and endometriosis-associated ovarian cancer. *Clin. Cancer Res.*, 20:6163-6174.
26. Măluțan, A.M., Drugan, T., Ciortea, R., Mocan-Hognogi, R.F., Bucuri, C., Rada, M.P., Miha, D. 2015. Serum anti-inflammatory cytokines for the evaluation of inflammatory status in endometriosis. *J. Res. Med. Sci.*, 20:668–674.
27. Weisheng, B., Nezhat, C.H., Huang, G.F. et al. 2019. Discovering endometriosis biomarkers with multiplex cytokine arrays. *Clin. Proteom.*, 16, 28, <https://doi.org/10.1186/s12014-019-9248-y>
28. Xu, X., Mei, J., Zhang, B., Jiang, X.Y., Wang, L., Zhang, A.X., et al. 2025. Association between circulating cytokines and endometriosis: a mendelian randomization study. *J. Cell. Mol. Med.*, (7):e70532. doi: 10.1111/jcmm.70532.
29. Krygere, L., Jukna, P., Jariene, K., Drejeriene, E. 2024. Diagnostic potential of cytokine biomarkers in endometriosis: Challenges and insights. *Biomedicines*, 12:2867, <https://doi.org/10.3390/biomedicines12122867>
30. Li, W., Lin, A., Qi, L., Lv, X., Yan, S., Xue, J., Mu, N. 2023. Immunotherapy: A promising novel endometriosis therapy. *Frontiers Immunol.*, 14:1128301, doi: 10.3389/fimmu.2023.1128301
31. Garza, D., Mathur, S., Dowd, M.M., Smith, L.F., Williamson, D.Q. 1991. Antigenic differences between the endometrium of women with and without endometriosis. *J. Reprod. Med.*, 26:177-182.
32. Pasoto, S.G., Abrao, M.S., Viana, V.S., Bueno, C., Leon, E.P., Bonfa, E. 2005. Endometriosis and systemic lupus erythematosus: a comparative evaluation of clinical manifestations and serological autoimmune phenomena. *Am. J. Reprod. Immunol.*, 53:85-93.
33. Zhang, T., De Carolis, C., Man, G.C.W., Wang, C.C. 2018. The link between the immunity, autoimmunity and endometriosis: a literature update. *Autoimmune Rev.*, 17:945-955.
34. Melioli, G., Semino, C., Semino, A., Venturini, P.L., Ragni, N. 1993. Recombinant interleukin-2 corrects in vitro the immunological defect of endometriosis. *Am. J. Reprod. Immunol.*, 30:218-227.
35. Czubak, P., Herda, K., Niewiadomska, I., Putowski, L., Łańcut, M., Małyk, M. 2025. Understanding endometriosis: A broad review of its causes, management, and impact. *Int. J. Mol. Sci.*, 26, 8878. Doi:10.3390/ijms26188878